

SCF (KITLG) Antibody (N-term) Blocking peptide

Synthetic peptide Catalog # BP1484a

Specification

SCF (KITLG) Antibody (N-term) Blocking peptide - Product Information

Primary Accession P21583

Other Accession Q28132, P79368

SCF (KITLG) Antibody (N-term) Blocking peptide - Additional Information

Gene ID 4254

Other Names

Kit ligand, Mast cell growth factor, MGF, Stem cell factor, SCF, c-Kit ligand, Soluble KIT ligand, sKITLG, KITLG, MGF, SCF

Target/Specificity

The synthetic peptide sequence is selected from aa 38-52 of HUMAN KITLG

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

SCF (KITLG) Antibody (N-term) Blocking peptide - Protein Information

Name KITLG (HGNC:6343)

Synonyms MGF, SCF

Function

Ligand for the receptor-type protein-tyrosine kinase KIT. Plays an essential role in the regulation of cell survival and proliferation, hematopoiesis, stem cell maintenance, gametogenesis, mast cell development, migration and function, and in melanogenesis. KITLG/SCF binding can activate several signaling pathways. Promotes phosphorylation of PIK3R1, the regulatory subunit of phosphatidylinositol 3-kinase, and subsequent activation of the kinase AKT1. KITLG/SCF and KIT also transmit signals via GRB2 and activation of RAS, RAF1 and the MAP kinases MAPK1/ERK2 and/or MAPK3/ERK1. KITLG/SCF and KIT promote activation of STAT family members STAT1, STAT3 and STAT5. KITLG/SCF and KIT promote activation of PLCG1, leading to the production of the cellular signaling molecules diacylglycerol and inositol 1,4,5-trisphosphate. KITLG/SCF acts synergistically with other cytokines, probably interleukins.



Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein [Soluble KIT ligand]: Secreted.

SCF (KITLG) Antibody (N-term) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

Blocking Peptides

SCF (KITLG) Antibody (N-term) Blocking peptide - Images

SCF (KITLG) Antibody (N-term) Blocking peptide - Background

KITLG is the ligand of the tyrosine-kinase receptor encoded by the KIT locus. This ligand is a pleiotropic factor that acts in utero in germ cell and neural cell development, and hematopoiesis, all believed to reflect a role in cell migration. In adults, it functions pleiotropically, while mostly noted for its continued requirement in hematopoiesis.

SCF (KITLG) Antibody (N-term) Blocking peptide - References

Young, S.M., Cell. Signal. 19 (12), 2572-2581 (2007) Pick, M., Stem Cells 25 (9), 2206-2214 (2007) Yasuda, A., Dig. Dis. Sci. 52 (9), 2292-2300 (2007)