

**SFRS3 Antibody (Center) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP16268c****Specification**

---

**SFRS3 Antibody (Center) Blocking Peptide - Product Information**Primary Accession [P84103](#)**SFRS3 Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 6428**Other Names**

Serine/arginine-rich splicing factor 3, Pre-mRNA-splicing factor SRP20, Splicing factor, arginine/serine-rich 3, SRSF3, SFRS3, SRP20

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**SFRS3 Antibody (Center) Blocking Peptide - Protein Information****Name** SRSF3**Synonyms** SFRS3, SRP20**Function**

Splicing factor, which binds the consensus motif 5'- C[ACU][AU]C[ACU][AC]C-3' within pre-mRNA and promotes specific exons inclusion during alternative splicing (PubMed:<a href="http://www.uniprot.org/citations/32440474" target="\_blank">32440474</a>, PubMed:<a href="http://www.uniprot.org/citations/17036044" target="\_blank">17036044</a>, PubMed:<a href="http://www.uniprot.org/citations/26876937" target="\_blank">26876937</a>). Interaction with YTHDC1, a RNA- binding protein that recognizes and binds N6-methyladenosine (m6A)-containing RNAs, promotes recruitment of SRSF3 to its mRNA-binding elements adjacent to m6A sites within exons (PubMed:<a href="http://www.uniprot.org/citations/26876937" target="\_blank">26876937</a>). Also functions as an adapter involved in mRNA nuclear export (PubMed:<a href="http://www.uniprot.org/citations/11336712" target="\_blank">11336712</a>, PubMed:<a href="http://www.uniprot.org/citations/18364396" target="\_blank">18364396</a>, PubMed:<a href="http://www.uniprot.org/citations/28984244" target="\_blank">28984244</a>). Binds mRNA which is thought to be transferred to the NXF1-NXT1 heterodimer for export (TAP/NXF1 pathway); enhances NXF1-NXT1 RNA-binding activity (PubMed:<a href="http://www.uniprot.org/citations/11336712" target="\_blank">11336712</a>, PubMed:<a

[18364396](http://www.uniprot.org/citations/18364396)). Involved in nuclear export of m6A- containing mRNAs via interaction with YTHDC1: interaction with YTHDC1 facilitates m6A-containing mRNA-binding to both SRSF3 and NXF1, promoting mRNA nuclear export (PubMed: [28984244](http://www.uniprot.org/citations/28984244)).

#### **Cellular Location**

Nucleus. Nucleus speckle. Cytoplasm. Note=Recruited to nuclear speckles following interaction with YTHDC1.

### **SFRS3 Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **SFRS3 Antibody (Center) Blocking Peptide - Images**

### **SFRS3 Antibody (Center) Blocking Peptide - Background**

SFRS3 is a member of the serine/arginine (SR)-rich family of pre-mRNA splicing factors, which constitute part of the spliceosome. Each of these factors contains an RNA recognition motif (RRM) for binding RNA and an RS domain for binding other proteins. The RS domain is rich in serine and arginine residues and facilitates interaction between different SR splicing factors. In addition to being critical for mRNA splicing, the SR proteins have also been shown to be involved in mRNA export from the nucleus and in translation. Two transcript variants, one protein-coding and the other non-coding, have been found for this gene.

### **SFRS3 Antibody (Center) Blocking Peptide - References**

Verma, D., et al. J. Virol. 84(22):11781-11789(2010) Anko, M.L., et al. Nat. Struct. Mol. Biol. 17(8):962-970(2010) Escudero-Paunetto, L., et al. Virology 401(2):155-164(2010) Manley, J.L., et al. Genes Dev. 24(11):1073-1074(2010) Fingert, J.H., et al. Mol. Vis. 16, 596-601 (2010) :