

IFT57 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP16420a**Specification**

IFT57 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [Q9NWB7](#)**IFT57 Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 55081**Other Names**

Intraflagellar transport protein 57 homolog, Dermal papilla-derived protein 8, Estrogen-related receptor beta-like protein 1, HIP1-interacting protein, MHS4R2, IFT57, DERP8, ESRRBL1, HIPPI

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

IFT57 Antibody (N-term) Blocking Peptide - Protein Information**Name** IFT57**Synonyms** DERP8, ESRRBL1, HIPPI**Function**

Required for the formation of cilia. Plays an indirect role in sonic hedgehog signaling, cilia being required for all activity of the hedgehog pathway (By similarity). Has pro-apoptotic function via its interaction with HIP1, leading to recruit caspase-8 (CASP8) and trigger apoptosis. Has the ability to bind DNA sequence motif 5'- AAAGACATG-3' present in the promoter of caspase genes such as CASP1, CASP8 and CASP10, suggesting that it may act as a transcription regulator; however the relevance of such function remains unclear.

Cellular Location

Cell projection, cilium {ECO:0000250|UniProtKB:Q8BXG3}. Cytoplasm, cytoskeleton, cilium basal body {ECO:0000250|UniProtKB:Q5EA95}. Note=Concentrates within the inner segment of cilia.

Tissue Location

Present in many tissues such as brain, thymus, lymph node, lung, liver, skin and kidney (at protein level)

IFT57 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

IFT57 Antibody (N-term) Blocking Peptide - Images

IFT57 Antibody (N-term) Blocking Peptide - Background

IFT57 is required for the formation of cilia. Plays an indirect role in sonic hedgehog signaling, cilia being required for all activity of the hedgehog pathway (By similarity). Has pro-apoptotic function via its interaction with HIP1, leading to recruit caspase-8 (CASP8) and trigger apoptosis. Has the ability to bind DNA sequence motif 5'-AAAGACATG-3' present in the promoter of caspase genes such as CASP1, CASP8 and CASP10, suggesting that it may act as a transcription regulator; however the relevance of such function remains unclear.

IFT57 Antibody (N-term) Blocking Peptide - References

Gdynia, G., et al. Apoptosis 13(3):437-447(2008)Majumder, P., et al. Biochem. Biophys. Res. Commun. 353(1):80-85(2007)Banerjee, M., et al. Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun. 62 (PT 12), 1247-1250 (2006) :Majumder, P., et al. Neurobiol. Dis. 22(2):242-256(2006)Fan, Y., et al. Nat. Genet. 36(9):989-993(2004)