

**THRAP3 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP16468b****Specification**

---

**THRAP3 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [O9Y2W1](#)**THRAP3 Antibody (C-term) Blocking Peptide - Additional Information**

Gene ID 9967

**Other Names**

Thyroid hormone receptor-associated protein 3, Thyroid hormone receptor-associated protein complex 150 kDa component, Trap150, THRAP3, TRAP150

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**THRAP3 Antibody (C-term) Blocking Peptide - Protein Information**Name THRAP3 ([HGNC:22964](#))**Function**

Involved in pre-mRNA splicing. Remains associated with spliced mRNA after splicing which probably involves interactions with the exon junction complex (EJC). Can trigger mRNA decay which seems to be independent of nonsense-mediated decay involving premature stop codons (PTC) recognition. May be involved in nuclear mRNA decay. Involved in regulation of signal-induced alternative splicing. During splicing of PTPRC/CD45 is proposed to sequester phosphorylated SFPQ from PTPRC/CD45 pre-mRNA in resting T-cells. Involved in cyclin- D1/CCND1 mRNA stability probably by acting as component of the SNARP complex which associates with both the 3'end of the CCND1 gene and its mRNA. Involved in response to DNA damage. Is excluded from DNA damage sites in a manner that parallels transcription inhibition; the function may involve the SNARP complex. Initially thought to play a role in transcriptional coactivation through its association with the TRAP complex; however, it is not regarded as a stable Mediator complex subunit. Cooperatively with HELZ2, enhances the transcriptional activation mediated by PPARG, maybe through the stabilization of the PPARG binding to DNA in presence of ligand. May play a role in the terminal stage of adipocyte differentiation. Plays a role in the positive regulation of the circadian clock. Acts as a coactivator of the CLOCK-BMAL1 heterodimer and promotes its transcriptional activator activity and binding to circadian target genes (PubMed:<a href="http://www.uniprot.org/citations/24043798" target="\_blank">24043798</a>).

**Cellular Location**

Nucleus. Nucleus, nucleoplasm. Nucleus speckle

**Tissue Location**

Ubiquitous..

**THRAP3 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**THRAP3 Antibody (C-term) Blocking Peptide - Images****THRAP3 Antibody (C-term) Blocking Peptide - Background**

THRAP3 plays a role in transcriptional coactivation.

**THRAP3 Antibody (C-term) Blocking Peptide - References**

Ewing, R.M., et al. Mol. Syst. Biol. 3, 89 (2007) :Olsen, J.V., et al. Cell  
127(3):635-648(2006)Beausoleil, S.A., et al. Nat. Biotechnol. 24(10):1285-1292(2006)Nousiainen,  
M., et al. Proc. Natl. Acad. Sci. U.S.A. 103(14):5391-5396(2006)Jin, J., et al. Curr. Biol.  
14(16):1436-1450(2004)