

# AP3S2 Antibody (N-term) Blocking Peptide

Synthetic peptide Catalog # BP16494a

## **Specification**

## AP3S2 Antibody (N-term) Blocking Peptide - Product Information

**Primary Accession** 

P59780

## AP3S2 Antibody (N-term) Blocking Peptide - Additional Information

**Gene ID** 10239

#### **Other Names**

AP-3 complex subunit sigma-2, AP-3 complex subunit sigma-3B, Adaptor-related protein complex 3 subunit sigma-2, Sigma-3B-adaptin, Sigma-adaptin, Sigma-adaptin 3b, AP3S2

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### AP3S2 Antibody (N-term) Blocking Peptide - Protein Information

#### Name AP3S2

## **Function**

Part of the AP-3 complex, an adaptor-related complex which is not clathrin-associated. The complex is associated with the Golgi region as well as more peripheral structures. It facilitates the budding of vesicles from the Golgi membrane and may be directly involved in trafficking to lysosomes. In concert with the BLOC-1 complex, AP-3 is required to target cargos into vesicles assembled at cell bodies for delivery into neurites and nerve terminals.

#### **Cellular Location**

Golgi apparatus. Cytoplasmic vesicle membrane; Peripheral membrane protein; Cytoplasmic side. Note=Component of the coat surrounding the cytoplasmic face of coated vesicles located at the Golgi complex

#### **Tissue Location**

Present in all adult tissues examined.

## AP3S2 Antibody (N-term) Blocking Peptide - Protocols



Provided below are standard protocols that you may find useful for product applications.

### • Blocking Peptides

## AP3S2 Antibody (N-term) Blocking Peptide - Images

# AP3S2 Antibody (N-term) Blocking Peptide - Background

Part of the AP-3 complex, an adapter-related complex which is not clathrin-associated. The complex is associated with the Golgi region as well as more peripheral structures. It facilitates the budding of vesicles from the Golgi membrane and may be directly involved in trafficking to lysosomes.

## AP3S2 Antibody (N-term) Blocking Peptide - References

Lefrancois, S., et al. Dev. Cell 7(4):619-625(2004)Nie, Z., et al. Dev. Cell 5(3):513-521(2003)Falcon-Perez, J.M., et al. J. Biol. Chem. 277(31):28191-28199(2002)Rous, B.A., et al. Mol. Biol. Cell 13(3):1071-1082(2002)Dell'Angelica, E.C., et al. J. Biol. Chem. 272(24):15078-15084(1997)