

ACSL3 Antibody (C-term) Blocking Peptide

Synthetic peptide Catalog # BP16647b

Specification

ACSL3 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession

095573

ACSL3 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 2181

Other Names

Long-chain-fatty-acid--CoA ligase 3, Long-chain acyl-CoA synthetase 3, LACS 3, ACSL3, ACSS, FACL3, LACS3

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ACSL3 Antibody (C-term) Blocking Peptide - Protein Information

Name ACSL3 (HGNC:3570)

Synonyms ACS3, FACL3, LACS3

Function

Acyl-CoA synthetases (ACSL) activates long-chain fatty acids for both synthesis of cellular lipids, and degradation via beta- oxidation (PubMed:22633490). Required for the incorporation of fatty acids into phosphatidylcholine, the major phospholipid located on the surface of VLDL (very low density lipoproteins) (PubMed:18003621). Has mainly an anabolic role in energy metabolism. Mediates hepatic lipogenesis. Preferentially uses myristate, laurate, arachidonate and eicosapentaenoate as substrates. Both isoforms exhibit the same level of activity (By similarity).

Cellular Location

Mitochondrion outer membrane; Single-pass type III membrane protein. Peroxisome membrane; Single-pass type III membrane protein. Microsome membrane; Single-pass type III membrane protein. Endoplasmic reticulum membrane; Single-pass type III membrane protein



ACSL3 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

ACSL3 Antibody (C-term) Blocking Peptide - Images

ACSL3 Antibody (C-term) Blocking Peptide - Background

The protein encoded by this gene is an isozyme of thelong-chain fatty-acid-coenzyme A ligase family. Although differingin substrate specificity, subcellular localization, and tissuedistribution, all isozymes of this family convert free long-chainfatty acids into fatty acyl-CoA esters, and thereby play a key rolein lipid biosynthesis and fatty acid degradation. This isozyme ishighly expressed in brain, and preferentially utilizes myristate, arachidonate, and eicosapentaenoate as substrates. The amino acidsequence of this isozyme is 92% identical to that of rat homolog. Two transcript variants encoding the same protein have been foundfor this gene.

ACSL3 Antibody (C-term) Blocking Peptide - References

Weedon-Fekjaer, M.S., et al. J. Lipid Res. 51(7):1886-1896(2010)Cao, A., et al. J. Biol. Chem. 285(22):16664-16674(2010)Perera, F., et al. PLoS ONE 4 (2), E4488 (2009) :Yao, H., et al. J. Biol. Chem. 283(2):849-854(2008)Jia, Z., et al. J. Mol. Neurosci. 33(1):25-31(2007)