

TRPC4 Antibody (C-term) Blocking Peptide
Synthetic peptide
Catalog # BP16760b**Specification**

TRPC4 Antibody (C-term) Blocking Peptide - Product Information

Primary Accession [Q9UBN4](#)

TRPC4 Antibody (C-term) Blocking Peptide - Additional Information

Gene ID 7223

Other Names

Short transient receptor potential channel 4, TrpC4, Trp-related protein 4, hTrp-4, hTrp4, TRPC4

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

TRPC4 Antibody (C-term) Blocking Peptide - Protein Information

Name TRPC4

Function

Forms a receptor-activated non-selective calcium permeant cation channel. Acts as a cell-cell contact-dependent endothelial calcium entry channel. Probably operated by a phosphatidylinositol second messenger system activated by receptor tyrosine kinases or G- protein coupled receptors. Mediates cation entry, with an enhanced permeability to barium over calcium. May also be activated by intracellular calcium store depletion.

Cellular Location

Membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein.
Note=Enhanced insertion into the cell membrane after activation of the EGF receptor

Tissue Location

Strongly expressed in placenta. Expressed at lower levels in heart, pancreas, kidney and brain. Expressed in endothelial cells. Isoform alpha was found to be the predominant isoform. Isoform beta was not found in pancreas and brain

TRPC4 Antibody (C-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

TRPC4 Antibody (C-term) Blocking Peptide - Images

TRPC4 Antibody (C-term) Blocking Peptide - Background

Form a receptor-activated non-selective calcium permeant cation channel. Acts as a cell-cell contact-dependent endothelial calcium entry channel. Probably operated by a phosphatidylinositol second messenger system activated by receptor tyrosine kinases or G-protein coupled receptors. Mediates cation entry, with an enhanced permeability to barium over calcium. May also be activated by intracellular calcium store depletion.

TRPC4 Antibody (C-term) Blocking Peptide - References

Mace, K.E., et al. J. Cell. Physiol. 225(2):444-453(2010) von Spiczak, S., et al. Neuromolecular Med. 12(3):292-299(2010) Pinheiro, A.P., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 153B (5), 1070-1080 (2010) :Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :Schaefer, M., et al. J. Biol. Chem. 277(5):3752-3759(2002)