

**FAM3C Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP16888b****Specification**

---

**FAM3C Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [Q92520](#)**FAM3C Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 10447**Other Names**

Protein FAM3C, Interleukin-like EMT inducer, FAM3C, ILEI

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**FAM3C Antibody (C-term) Blocking Peptide - Protein Information****Name** FAM3C**Synonyms** ILEI**Function**

May be involved in retinal laminar formation. Promotes epithelial to mesenchymal transition.

**Cellular Location**

Secreted. Cytoplasmic vesicle. Note=Cytoplasmic in some cancer cells

**Tissue Location**

Present in most secretory epithelia (at protein level).

**FAM3C Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**FAM3C Antibody (C-term) Blocking Peptide - Images**

**FAM3C Antibody (C-term) Blocking Peptide - Background**

This gene is a member of the family with sequence similarity 3 (FAM3) family and encodes a secreted protein with a GG domain. A change in expression of this protein has been noted in pancreatic cancer-derived cells.

**FAM3C Antibody (C-term) Blocking Peptide - References**

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :Cho, Y.S., et al. Nat. Genet. 41(5):527-534(2009)Xiao, F., et al. Brain Res. 1255, 180-189 (2009) :Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007)Waerner, T., et al. Cancer Cell 10(3):227-239(2006)