

## MRP63 Antibody (Center) Blocking Peptide

Synthetic peptide Catalog # BP16939c

### **Specification**

## MRP63 Antibody (Center) Blocking Peptide - Product Information

**Primary Accession** 

Q9BQC6

# MRP63 Antibody (Center) Blocking Peptide - Additional Information

**Gene ID** 78988

#### **Other Names**

Ribosomal protein 63, mitochondrial, hMRP63, Mitochondrial ribosomal protein 63, Mitochondrial ribosomal protein L57, MRPL57, MRP63

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### MRP63 Antibody (Center) Blocking Peptide - Protein Information

Name MRPL57

Synonyms MRP63

### **Cellular Location**

Mitochondrion

# MRP63 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

### • Blocking Peptides

MRP63 Antibody (Center) Blocking Peptide - Images

### MRP63 Antibody (Center) Blocking Peptide - Background

Mammalian mitochondrial ribosomal proteins are encoded bynuclear genes and help in protein synthesis within themitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of asmall 28S





Tel: 858.875.1900 Fax: 858.875.1999

subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalianmitoribosomes and prokaryotic ribosomes is that the latter containa 5S rRNA. Among different species, the proteins comprising themitoribosome differ greatly in sequence, and sometimes inbiochemical properties, which prevents easy recognition by sequencehomology. This gene encodes a protein which belongs to anundetermined ribosomal subunit and which seems to be specific toanimal mitoribosomes. Pseudogenes corresponding to this gene arefound on chromosomes 1p, 1q, 3p, 5q, 8q, 14q, and Y. [provided byRefSeq].

### MRP63 Antibody (Center) Blocking Peptide - References

Zhang, Z., et al. Genomics 81(5):468-480(2003)Kenmochi, N., et al. Genomics 77 (1-2), 65-70 (2001) :Suzuki, T., et al. J. Biol. Chem. 276(35):33181-33195(2001)