

NDOR1 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP17396c**Specification**

NDOR1 Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [Q9UHB4](#)**NDOR1 Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 27158**Other Names**

NADPH-dependent diflavin oxidoreductase 1 {ECO:0000255|HAMAP-Rule:MF_03178}, 16--
{ECO:0000255|HAMAP-Rule:MF_03178}, NADPH-dependent FMN and FAD-containing
oxidoreductase {ECO:0000255|HAMAP-Rule:MF_03178}, Novel reductase 1, NDOR1
{ECO:0000255|HAMAP-Rule:MF_03178}, NR1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

NDOR1 Antibody (Center) Blocking Peptide - Protein Information**Name** NDOR1 {ECO:0000255|HAMAP-Rule:MF_03178}**Function**

NADPH-dependent reductase which is a central component of the cytosolic iron-sulfur (Fe-S) protein assembly (CIA) machinery (PubMed: 10625700, PubMed: 28648056, PubMed: 23596212, PubMed: 20802492, PubMed: 15900210). Transfers electrons from NADPH via its FAD and FMN prosthetic groups to the [2Fe-2S] cluster of CIAPIN1, another key component of the CIA machinery (PubMed: 28648056, PubMed: 23596212, PubMed: 20802492). In turn, this reduced cluster provides electrons for assembly of cytosolic iron-sulfur cluster proteins (PubMed: 23596212, PubMed: 20802492). It can also reduce the [2Fe-2S] cluster of CISD1 and activate this protein implicated in Fe/S cluster

repair (PubMed:28648056). In vitro can fully activate methionine synthase/MTR in the presence of soluble cytochrome b5/CYB5A (PubMed:12871938).

Cellular Location

Cytoplasm, perinuclear region {ECO:0000255|HAMAP- Rule:MF_03178, ECO:0000269|PubMed:10625700, ECO:0000269|PubMed:12871939}. Note=Concentrated in perinuclear structure. {ECO:0000255|HAMAP-Rule:MF_03178, ECO:0000269|PubMed:12871939}

Tissue Location

Low expression in brain, heart, kidney, pancreas, prostate and skeletal muscle. Highest levels in the placenta. Expressed in cancer cell lines including promyelocytic leukemia, HeLaS3, chronic myelogenous leukemia, lymphoblastic leukemia, Burkitt's lymphoma, colorectal adenocarcinoma, lung carcinoma, and melanoma G-361

NDOR1 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

NDOR1 Antibody (Center) Blocking Peptide - Images

NDOR1 Antibody (Center) Blocking Peptide - Background

This gene encodes an NADPH-dependent diflavin reductase that contains both flavin mononucleotide (FMN) and flavin adenine dinucleotide (FAD) binding domains. The encoded protein is an enzyme that catalyzes the transfers electrons from NADPH through FAD and FMN cofactors to potential redox partners. Alternative splicing results in multiple transcript variants. [provided by RefSeq].

NDOR1 Antibody (Center) Blocking Peptide - References

Kwasnicka-Crawford, D.A., et al. Biochem. Biophys. Res. Commun. 336(2):565-571(2005) Finn, R.D., et al. Pharmacogenet. Genomics 15(6):381-386(2005) Kwasnicka, D.A., et al. J. Biol. Chem. 278(40):39051-39058(2003) Olteanu, H., et al. J. Biol. Chem. 278(40):38310-38314(2003) Finn, R.D., et al. Eur. J. Biochem. 270(6):1164-1175(2003)