

WIPI1 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP17541c**Specification**

WIPI1 Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [Q5MNZ9](#)**WIPI1 Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 55062**Other Names**

WD repeat domain phosphoinositide-interacting protein 1, WIPI-1, Atg18 protein homolog, WD40 repeat protein interacting with phosphoinositides of 49 kDa, WIPI 49 kDa, WIPI1, WIPI49

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

WIPI1 Antibody (Center) Blocking Peptide - Protein Information**Name** WIPI1**Synonyms** WIPI49**Function**

Component of the autophagy machinery that controls the major intracellular degradation process by which cytoplasmic materials are packaged into autophagosomes and delivered to lysosomes for degradation (PubMed: [15602573](http://www.uniprot.org/citations/15602573), PubMed: [20114074](http://www.uniprot.org/citations/20114074), PubMed: [20484055](http://www.uniprot.org/citations/20484055), PubMed: [20639694](http://www.uniprot.org/citations/20639694), PubMed: [23088497](http://www.uniprot.org/citations/23088497), PubMed: [28561066](http://www.uniprot.org/citations/28561066), PubMed: [31271352](http://www.uniprot.org/citations/31271352)). Plays an important role in starvation- and calcium-mediated autophagy, as well as in mitophagy (PubMed: [28561066](http://www.uniprot.org/citations/28561066)). Functions downstream of the ULK1 and PI3- kinases that produce phosphatidylinositol 3-phosphate (PtdIns3P) on membranes of the endoplasmic reticulum once activated (PubMed: [28561066](http://www.uniprot.org/citations/28561066)). Binds phosphatidylinositol 3-phosphate (PtdIns3P), and maybe

other phosphoinositides including PtdIns3,5P2 and PtdIns5P, and is recruited to phagophore assembly sites at the endoplasmic reticulum membranes (PubMed:28561066, PubMed:31271352, PubMed:33499712). There, it assists WIPI2 in the recruitment of ATG12- ATG5-ATG16L1, a complex that directly controls the elongation of the nascent autophagosomal membrane (PubMed:28561066). Together with WDR45/WIPI4, promotes ATG2 (ATG2A or ATG2B)-mediated lipid transfer by enhancing ATG2-association with phosphatidylinositol 3-monophosphate (PI3P)-containing membranes (PubMed:31271352). Involved in xenophagy of Staphylococcus aureus (PubMed:22829830). Invading S.aureus cells become entrapped in autophagosome-like WIPI1 positive vesicles targeted for lysosomal degradation (PubMed:22829830). Also plays a distinct role in controlling the transcription of melanogenic enzymes and melanosome maturation, a process that is distinct from starvation-induced autophagy (PubMed:21317285). May also regulate the trafficking of proteins involved in the mannose-6-phosphate receptor (MPR) recycling pathway (PubMed:15020712).

Cellular Location

Golgi apparatus, trans-Golgi network. Endosome. Cytoplasmic vesicle, clathrin-coated vesicle. Preautophagosomal structure membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Note=Trans elements of the Golgi and peripheral endosomes. Dynamically cycles through these compartments and is susceptible to conditions that modulate membrane flux. Enriched in clathrin-coated vesicles. Upon starvation-induced autophagy, accumulates at subcellular structures in the cytoplasm: enlarged vesicular and lasso-like structures, and large cup-shaped structures predominantly around the nucleus. Recruitment to autophagic membranes is controlled by MTMR14. Labile microtubules specifically recruit markers of autophagosome formation like WIPI1, whereas mature autophagosomes may bind to stable microtubules

Tissue Location

Ubiquitously expressed. Highly expressed in skeletal muscle, heart, testis, pancreas and placenta. Highly expressed in G361, Sk-mel-28, Sk-mel-13, WM852 and WM451 cells. Up-regulated in a variety of tumor tissues.

WIPI1 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

WIPI1 Antibody (Center) Blocking Peptide - Images

WIPI1 Antibody (Center) Blocking Peptide - Background

WD40 repeat proteins are key components of many essentialbiologic functions. They regulate the assembly of multiproteincomplexes by presenting a beta-propeller platform for simultaneousand reversible protein-protein interactions. Members of the WIPIsubfamily of WD40 repeat proteins, such as WIPI1, have a 7-bladedpropeller structure and contain a conserved motif for interactionwith phospholipids (Proikas-Cezanne et al., 2004 [PubMed15602573]).

WIPI1 Antibody (Center) Blocking Peptide - References

Chasman, D.I., et al. PLoS Genet. 5 (11), E1000730 (2009) :Proikas-Cezanne, T., et al. FEBS Lett. 581(18):3396-3404(2007)Proikas-Cezanne, T., et al. Oncogene 23(58):9314-9325(2004)Jeffries, T.R., et al. Mol. Biol. Cell 15(6):2652-2663(2004)