

## LPHN3 Antibody (N-term) Blocking Peptide

Synthetic peptide Catalog # BP17661a

## **Specification**

## LPHN3 Antibody (N-term) Blocking Peptide - Product Information

**Primary Accession** 

Q9HAR2

## LPHN3 Antibody (N-term) Blocking Peptide - Additional Information

**Gene ID 23284** 

#### **Other Names**

Latrophilin-3, Calcium-independent alpha-latrotoxin receptor 3, CIRL-3, Lectomedin-3, LPHN3, KIAA0768, LEC3

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

## **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## LPHN3 Antibody (N-term) Blocking Peptide - Protein Information

Name ADGRL3 (HGNC:20974)

## **Function**

Plays a role in cell-cell adhesion and neuron guidance via its interactions with FLRT2 and FLRT3 that are expressed at the surface of adjacent cells (PubMed:<a href="http://www.uniprot.org/citations/26235030" target="\_blank">26235030</a>). Plays a role in the development of glutamatergic synapses in the cortex. Important in determining the

#### **Cellular Location**

Cell membrane; Multi-pass membrane protein. Cell projection, axon {ECO:0000250|UniProtKB:Q80TS3}. Cell junction {ECO:0000250|UniProtKB:Q80TS3}

### LPHN3 Antibody (N-term) Blocking Peptide - Protocols

connectivity rates between the principal neurons in the cortex.

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides



Tel: 858.875.1900 Fax: 858.875.1999



# LPHN3 Antibody (N-term) Blocking Peptide - Images LPHN3 Antibody (N-term) Blocking Peptide - Background

This gene encodes a member of the latrophilin subfamily of G-protein coupled receptors (GPCR). Latrophilins may function inboth cell adhesion and signal transduction. In experiments withnon-human species, endogenous proteolytic cleavage within acysteine-rich GPS (G-protein-coupled-receptor proteolysis site)domain resulted in two subunits (a large extracellular N-terminalcell adhesion subunit and a subunit with substantial similarity tothe secretin/calcitonin family of GPCRs) being non-covalently boundat the cell membrane.

## LPHN3 Antibody (N-term) Blocking Peptide - References

Arcos-Burgos, M., et al. Mol. Psychiatry 15(11):1053-1066(2010)Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)Kasperaviciute, D., et al. Brain 133 (PT 7), 2136-2147 (2010):Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :Zemunik, T., et al. Croat. Med. J. 50(1):23-33(2009)