

**SLC12A4 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP18058b****Specification**

---

**SLC12A4 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [Q9UP95](#)**SLC12A4 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 6560**Other Names**

Solute carrier family 12 member 4, Electroneutral potassium-chloride cotransporter 1, Erythroid K-Cl cotransporter 1, hKCC1, SLC12A4, KCC1

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**SLC12A4 Antibody (C-term) Blocking Peptide - Protein Information****Name** SLC12A4 ([HGNC:10913](#))**Function**

Mediates electroneutral potassium-chloride cotransport when activated by cell swelling. May contribute to cell volume homeostasis in single cells (PubMed:&lt;a href="http://www.uniprot.org/citations/10913127" target="\_blank"&gt;10913127&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/34031912" target="\_blank"&gt;34031912&lt;/a&gt;). May be involved in the regulation of basolateral Cl(-) exit in NaCl absorbing epithelia (By similarity).

**Cellular Location**

Cell membrane; Multi-pass membrane protein

**Tissue Location**

Ubiquitous (PubMed:8663127, PubMed:9516379). Levels are much higher in erythrocytes from patients with Hb SC and Hb SS compared to normal AA erythrocytes (PubMed:9516379). This may contribute to red blood cell dehydration and to the manifestation of sickle cell disease by increasing the intracellular concentration of HbS (PubMed:9516379).

**SLC12A4 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**SLC12A4 Antibody (C-term) Blocking Peptide - Images****SLC12A4 Antibody (C-term) Blocking Peptide - Background**

SLC12A4 mediates electroneutral potassium-chloride cotransport when activated by cell swelling. May contribute to cell volume homeostasis in single cells. May be involved in the regulation of basolateral Cl(-) exit in NaCl absorbing epithelia (By similarity). Isoform 4 has no transport activity.

**SLC12A4 Antibody (C-term) Blocking Peptide - References**

Bailey, S.D., et al. Diabetes Care (2010) In press :Ruano, G., et al. Pharmacogenomics 11(7):959-971(2010)Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)Zhang, S., et al. Eur. J. Gynaecol. Oncol. 30(1):29-34(2009)Zhou, G.P., et al. Blood 103(11):4302-4309(2004)