

## **ELL3 Antibody (C-term) Blocking Peptide**

Synthetic peptide Catalog # BP18266b

## **Specification**

# **ELL3 Antibody (C-term) Blocking Peptide - Product Information**

**Primary Accession** 

**Q9HB65** 

# ELL3 Antibody (C-term) Blocking Peptide - Additional Information

**Gene ID 80237** 

#### **Other Names**

RNA polymerase II elongation factor ELL3, ELL3

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### ELL3 Antibody (C-term) Blocking Peptide - Protein Information

## Name ELL3

### **Function**

Enhancer-binding elongation factor that specifically binds enhancers in embryonic stem cells (ES cells), marks them, and is required for their future activation during stem cell specification. Does not only bind to enhancer regions of active genes, but also marks the enhancers that are in a poised or inactive state in ES cells and is required for establishing proper RNA polymerase II occupancy at developmentally regulated genes in a cohesin-dependent manner. Probably required for priming developmentally regulated genes for later recruitment of the super elongation complex (SEC), for transcriptional activation during differentiation. Required for recruitment of P-TEFb within SEC during differentiation. Probably preloaded on germ cell chromatin, suggesting that it may prime gene activation by marking enhancers as early as in the germ cells. Promoting epithelial- mesenchymal transition (EMT) (By similarity). Elongation factor component of the super elongation complex (SEC), a complex required to increase the catalytic rate of RNA polymerase II transcription by suppressing transient pausing by the polymerase at multiple sites along the DNA. Component of the little elongation complex (LEC), a complex required to regulate small nuclear RNA (snRNA) gene transcription by RNA polymerase II and III (PubMed:<a href="http://www.uniprot.org/citations/22195968" target="blank">22195968</a>/a>).

#### **Cellular Location**

Nucleus.



**Tissue Location** Testis specific..

# **ELL3 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

**ELL3 Antibody (C-term) Blocking Peptide - Images** 

# ELL3 Antibody (C-term) Blocking Peptide - Background

ELL3 is a elongation factor that can increase the catalytic rate of RNA polymerase II transcription by suppressing transient pausing by the polymerase at multiple sites along the DNA.

# ELL3 Antibody (C-term) Blocking Peptide - References

Lamesch, P., et al. Genomics 89(3):307-315(2007)Miller, T., et al. J. Biol. Chem. 275(41):32052-32056(2000)