

ATP5L2 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP18767a**Specification**

ATP5L2 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession [Q7Z4Y8](#)**ATP5L2 Antibody (N-term) Blocking Peptide - Additional Information****Gene ID** 267020**Other Names**

ATP synthase subunit g 2, mitochondrial, ATPase subunit g 2, ATP5L2, ATP5K2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

ATP5L2 Antibody (N-term) Blocking Peptide - Protein Information**Name** ATP5MGL ([HGNC:13213](#))**Synonyms** ATP5K2, ATP5L2**Function**

Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) - containing the extramembraneous catalytic core, and F(0) - containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F(0) domain. Minor subunit located with subunit a in the membrane (By similarity).

Cellular Location

Mitochondrion membrane.

ATP5L2 Antibody (N-term) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

ATP5L2 Antibody (N-term) Blocking Peptide - Images

ATP5L2 Antibody (N-term) Blocking Peptide - Background

Mitochondrial membrane ATP synthase (F(1)F(0) ATP synthase or Complex V) produces ATP from ADP in the presence of a proton gradient across the membrane which is generated by electron transport complexes of the respiratory chain. F-type ATPases consist of two structural domains, F(1) -containing the extramembraneous catalytic core, and F(0) -containing the membrane proton channel, linked together by a central stalk and a peripheral stalk. During catalysis, ATP synthesis in the catalytic domain of F(1) is coupled via a rotary mechanism of the central stalk subunits to proton translocation. Part of the complex F(0) domain. Minor subunit located with subunit a in the membrane (By similarity).

ATP5L2 Antibody (N-term) Blocking Peptide - References

Gerhard, D.S., et al. Genome Res. 14 (10B), 2121-2127 (2004) :