

### **RALGDS Antibody (Center) Blocking Peptide**

Synthetic peptide Catalog # BP18831c

### **Specification**

### **RALGDS Antibody (Center) Blocking Peptide - Product Information**

Primary Accession

**Q12967** 

# **RALGDS Antibody (Center) Blocking Peptide - Additional Information**

**Gene ID 5900** 

#### **Other Names**

Ral guanine nucleotide dissociation stimulator, RalGDS, Ral guanine nucleotide exchange factor, RalGEF, RALGDS, KIAA1308, RGF

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### **Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

### **RALGDS Antibody (Center) Blocking Peptide - Protein Information**

**Name RALGDS** 

Synonyms KIAA1308, RGF

#### **Function**

Functions as a guanine nucleotide exchange factor (GEF) activating either RalA or RalB GTPases and plays an important role in intracellular transport. Interacts and acts as an effector molecule for R-Ras, H-Ras, K-Ras, and Rap (By similarity). During bacterial clearance, recognizes 'Lys-33'-linked polyubiquitinated TRAF3 and subsequently mediates assembly of the exocyst complex (PubMed:<a href="http://www.uniprot.org/citations/27438768" target="\_blank">27438768</a>).

#### **Cellular Location**

Cytoplasm {ECO:0000250|UniProtKB:Q03385}. Nucleus {ECO:0000250|UniProtKB:Q03385}. Note=Localizes mainly in the peripheral region of the cytoplasmic membrane in oocytes and in preimplantation embryos until the 8-cell stage. Between the late 1-cell and the early 2-cell stages, nuclear localization becomes stronger. After the 4-cell stage, not detected in the nucleus. {ECO:0000250|UniProtKB:Q03385}



# **RALGDS Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

### • Blocking Peptides

**RALGDS Antibody (Center) Blocking Peptide - Images** 

#### RALGDS Antibody (Center) Blocking Peptide - Background

Guanine nucleotide dissociation stimulators (GDSs, orexchange factors), such as RALGDS, are effectors of Ras-relatedGTPases (see MIM 190020) that participate in signaling for avariety of cellular processes.

# **RALGDS Antibody (Center) Blocking Peptide - References**

Fuentes, G., et al. Trends Biochem. Sci. 34(11):533-539(2009)Maehama, T., et al. J. Biol. Chem. 283(50):35053-35059(2008)Rondaij, M.G., et al. Blood 112(1):56-63(2008)Omholt, K., et al. Melanoma Res. 17(6):410-412(2007)Yin, J., et al. Mol. Cell. Biol. 27(21):7538-7550(2007)