

SNK Antibody (Center K390) Blocking Peptide Synthetic peptide Catalog # BP18959c

Specification

SNK Antibody (Center K390) Blocking Peptide - Product Information

Primary Accession

<u>Q9NYY3</u>

SNK Antibody (Center K390) Blocking Peptide - Additional Information

Gene ID 10769

Other Names

Serine/threonine-protein kinase PLK2, Polo-like kinase 2, PLK-2, hPlk2, Serine/threonine-protein kinase SNK, hSNK, Serum-inducible kinase, PLK2, SNK

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions This product is for research use only. Not for use in diagnostic or therapeutic procedures.

SNK Antibody (Center K390) Blocking Peptide - Protein Information

Name PLK2

Synonyms SNK

Function

Tumor suppressor serine/threonine-protein kinase involved in synaptic plasticity, centriole duplication and G1/S phase transition. Polo-like kinases act by binding and phosphorylating proteins that are already phosphorylated on a specific motif recognized by the POLO box domains. Phosphorylates CPAP, NPM1, RAPGEF2, RASGRF1, SNCA, SIPA1L1 and SYNGAP1. Plays a key role in synaptic plasticity and memory by regulating the Ras and Rap protein signaling: required for overactivity-dependent spine remodeling by phosphorylating the Ras activator RASGRF1 and the Rap inhibitor SIPA1L1 leading to their degradation by the proteasome. Conversely, phosphorylates the Rap activator RAPGEF2 and the Ras inhibitor SYNGAP1, promoting their activity. Also regulates synaptic plasticity independently of kinase activity, via its interaction with NSF that disrupts the interaction between NSF and the GRIA2 subunit of AMPARs, leading to a rapid rundown of AMPAR-mediated current that occludes long term depression. Required for procentriole formation and centriole duplication by phosphorylating CPAP and NPM1, respectively. Its induction by p53/TP53 suggests that it may participate in the mitotic checkpoint following stress.

Cellular Location



Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cell projection, dendrite Note=Localizes to centrosomes during early G1 phase where it only associates to the mother centriole and then distributes equally to both mother and daughter centrioles at the onset of S phase

Tissue Location

Expressed at higher level in the fetal lung, kidney, spleen and heart.

SNK Antibody (Center K390) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

<u>Blocking Peptides</u>

SNK Antibody (Center K390) Blocking Peptide - Images

SNK Antibody (Center K390) Blocking Peptide - Background

Serum-inducible kinase is a member of the 'polo' family ofserine/threonine protein kinases that have a role in normal celldivision.

SNK Antibody (Center K390) Blocking Peptide - References

Chang, J., et al. EMBO J. 29(14):2395-2406(2010)Olson, J.E., et al. Breast Cancer Res. Treat. (2010) In press :Couch, F.J., et al. Cancer Epidemiol. Biomarkers Prev. 19(1):251-257(2010)Matthew, E.M., et al. Cell Cycle 8(24):4168-4175(2009)Wheeler, H.E., et al. PLoS Genet. 5 (10), E1000685 (2009) :