

UBAP2L Antibody (Center) Blocking Peptide

Synthetic peptide Catalog # BP19097c

Specification

UBAP2L Antibody (Center) Blocking Peptide - Product Information

Primary Accession

<u>014157</u>

UBAP2L Antibody (Center) Blocking Peptide - Additional Information

Gene ID 9898

Other Names

Ubiquitin-associated protein 2-like, Protein NICE-4, UBAP2L, KIAA0144, NICE4

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

UBAP2L Antibody (Center) Blocking Peptide - Protein Information

Name UBAP2L (HGNC:29877)

Synonyms KIAA0144, NICE4

Function

Recruits the ubiquitination machinery to RNA polymerase II for polyubiquitination, removal and degradation, when the transcription-coupled nucleotide excision repair (TC-NER) machinery fails to resolve DNA damage (PubMed:<a href="http://www.uniprot.org/citations/35633597"

target="_blank">35633597). Plays an important role in the activity of long-term repopulating hematopoietic stem cells (LT- HSCs) (By similarity). Is a regulator of stress granule assembly, required for their efficient formation (PubMed:<a

 $href="http://www.uniprot.org/citations/29395067" target="_blank">29395067, PubMed:35977029). Required for proper brain development and neocortex lamination (By similarity).$

Cellular Location

Nucleus. Chromosome. Cytoplasm Cytoplasm, Stress granule Note=Associates with nuclear chromatin.

Tissue Location

Ubiquitous..



UBAP2L Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

UBAP2L Antibody (Center) Blocking Peptide - Images

UBAP2L Antibody (Center) Blocking Peptide - Background

UBAP2L contains 1 UBA domain and has 3 named isoforms. UBA domains are a commonly occurring sequence motif of approximately 45 amino acid residues that are found in diverse proteins involved in the ubiquitin/proteasome pathway, DNA excision-repair, and cell signalling via protein kinases. The function of this protein is not known.

UBAP2L Antibody (Center) Blocking Peptide - References

Ichikawa, S., et al. J. Bone Miner. Res. 25(8):1821-1829(2010)Davila, S., et al. Genes Immun. 11(3):232-238(2010)Wang, Y.L., et al. J. Biol. Chem. 283(49):33808-33815(2008)Olsen, J.V., et al. Cell 127(3):635-648(2006)Beausoleil, S.A., et al. Nat. Biotechnol. 24(10):1285-1292(2006)