

**Mouse Mcm2 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP19218b****Specification**

---

**Mouse Mcm2 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [P97310](#)**Mouse Mcm2 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 17216**Other Names**

DNA replication licensing factor MCM2, Minichromosome maintenance protein 2 homolog, Nuclear protein BM28, Mcm2, Bm28, Cdc11, Kiaa0030, Mcmd2

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Mouse Mcm2 Antibody (C-term) Blocking Peptide - Protein Information****Name** Mcm2**Synonyms** Bm28, Cdc11, Kiaa0030, Mcmd2**Function**

Acts as a component of the MCM2-7 complex (MCM complex) which is the replicative helicase essential for 'once per cell cycle' DNA replication initiation and elongation in eukaryotic cells. Core component of CDC45-MCM-GINS (CMG) helicase, the molecular machine that unwinds template DNA during replication, and around which the replisome is built. The active ATPase sites in the MCM2-7 ring are formed through the interaction surfaces of two neighboring subunits such that a critical structure of a conserved arginine finger motif is provided in trans relative to the ATP-binding site of the Walker A box of the adjacent subunit. The six ATPase active sites, however, are likely to contribute differentially to the complex helicase activity. Required for the entry in S phase and for cell division (PubMed:<a href="http://www.uniprot.org/citations/10567526" target="\_blank">10567526</a>). Plays a role in terminally differentiated hair cells development of the cochlea and induces cells apoptosis (By similarity).

**Cellular Location**

Nucleus {ECO:0000250|UniProtKB:P49736}. Chromosome {ECO:0000250|UniProtKB:P49736}.  
Note=Associated with chromatin before the formation of nuclei and detaches from it as DNA

replication progresses. {ECO:0000250|UniProtKB:P55861}

### **Mouse Mcm2 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **Mouse Mcm2 Antibody (C-term) Blocking Peptide - Images**

### **Mouse Mcm2 Antibody (C-term) Blocking Peptide - Background**

Mcm2 acts as a factor that allows the DNA to undergo a single round of replication per cell cycle. Required for the entry in S phase and for cell division.

### **Mouse Mcm2 Antibody (C-term) Blocking Peptide - References**

Kunnev, D., et al. Oncogene 29(25):3630-3638(2010)Chuang, C.H., et al. PLoS Genet. 6 (9) (2010)  
:Lee, E.Y., et al. Dev. Biol. 332(1):104-115(2009)Hasegawa, M., et al. Leuk. Res.  
33(8):1100-1107(2009)Breunig, J.J., et al. Proc. Natl. Acad. Sci. U.S.A. 105(35):13127-13132(2008)