

**Mouse Dcc Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP19225b**

**Specification**

---

**Mouse Dcc Antibody (C-term) Blocking Peptide - Product Information**

Primary Accession [P70211](#)

**Mouse Dcc Antibody (C-term) Blocking Peptide - Additional Information**

**Gene ID** 13176

**Other Names**

Netrin receptor DCC, Tumor suppressor protein DCC, Dcc

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Mouse Dcc Antibody (C-term) Blocking Peptide - Protein Information**

**Name** Dcc

**Function**

Receptor for netrin required for axon guidance. Mediates axon attraction of neuronal growth cones in the developing nervous system upon ligand binding. Its association with UNC5 proteins may trigger signaling for axon repulsion. It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand. Implicated as a tumor suppressor gene.

**Cellular Location**

Membrane; Single-pass type I membrane protein.

**Tissue Location**

In the embryo, expressed at high levels in the developing brain and neural tube. In the embryo, expressed in developing neurons of the telencephalic cortical plate and in developing brainstem nuclei (PubMed:28250456). In adult, highly expressed in brain with very low levels found in testis, heart and thymus. Isoform C is expressed only in the embryo

**Mouse Dcc Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

#### **Mouse Dcc Antibody (C-term) Blocking Peptide - Images**

#### **Mouse Dcc Antibody (C-term) Blocking Peptide - Background**

Receptor for netrin required for axon guidance. Mediates axon attraction of neuronal growth cones in the developing nervous system upon ligand binding. Its association with UNC5 proteins may trigger signaling for axon repulsion. It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand. Implicated as a tumor suppressor gene.