

APG4B Antibody (Center G254) Blocking peptide
Synthetic peptide
Catalog # BP19349c**Specification**

APG4B Antibody (Center G254) Blocking peptide - Product InformationPrimary Accession [Q9Y4P1](#)**APG4B Antibody (Center G254) Blocking peptide - Additional Information**

Gene ID 23192

Other Names

Cysteine protease ATG4B, 3422-, AUT-like 1 cysteine endopeptidase, Autophagin-1, Autophagy-related cysteine endopeptidase 1, Autophagy-related protein 4 homolog B, hAPG4B, ATG4B, APG4B, AUTL1, KIAA0943

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

APG4B Antibody (Center G254) Blocking peptide - Protein Information**Name** ATG4B {ECO:0000303|PubMed:15187094, ECO:0000312|HGNC:HGNC:20790}**Function**

Cysteine protease that plays a key role in autophagy by mediating both proteolytic activation and delipidation of ATG8 family proteins (PubMed:15169837, PubMed:15187094, PubMed:17347651, PubMed:19322194, PubMed:21177865, PubMed:22302004, PubMed:26378241, PubMed:27527864, PubMed:28633005, PubMed:28821708, PubMed:29232556, PubMed:30076329, PubMed:30443548, PubMed:30661429). Required for

canonical autophagy (macroautophagy), non-canonical autophagy as well as for mitophagy (PubMed:33773106, PubMed:33909989). The protease activity is required for proteolytic activation of ATG8 family proteins: cleaves the C-terminal amino acid of ATG8 proteins MAP1LC3A, MAP1LC3B, MAP1LC3C, GABARAPL1, GABARAPL2 and GABARAP, to reveal a C- terminal glycine (PubMed:15169837, PubMed:15187094, PubMed:17347651, PubMed:19322194, PubMed:20818167, PubMed:21177865, PubMed:22302004, PubMed:27527864, PubMed:28287329, PubMed:28633005, PubMed:29458288, PubMed:30661429). Exposure of the glycine at the C-terminus is essential for ATG8 proteins conjugation to phosphatidylethanolamine (PE) and insertion to membranes, which is necessary for autophagy (PubMed:15169837, PubMed:15187094, PubMed:17347651, PubMed:19322194, PubMed:21177865, PubMed:22302004). Protease activity is also required to counteract formation of high-molecular weight conjugates of ATG8 proteins (ATG8ylation): acts as a deubiquitinating-like enzyme that removes ATG8 conjugated to other proteins, such as ATG3 (PubMed:31315929, PubMed:33773106). In addition to the protease activity, also mediates delipidation of ATG8 family proteins (PubMed:15187094, PubMed:19322194, PubMed:28633005, PubMed:29458288, PubMed:32686895, PubMed:33909989). Catalyzes delipidation of PE- conjugated forms of ATG8 proteins during macroautophagy (PubMed:15187094, PubMed:19322194, PubMed:29458288, PubMed:32686895, PubMed:33909989). Also involved in non-canonical autophagy, a parallel pathway involving conjugation of ATG8 proteins to single membranes at endolysosomal compartments, by catalyzing delipidation of ATG8 proteins conjugated to phosphatidylserine (PS) (PubMed:33909989). Compared to other members of the family (ATG4A, ATG4C or ATG4C), constitutes the major protein for proteolytic activation of ATG8 proteins, while it displays weaker delipidation activity than other ATG4 paralogs (PubMed:29458288, PubMed:30661429). Involved in phagophore growth during mitophagy independently of its protease activity and of ATG8 proteins: acts by regulating ATG9A trafficking to mitochondria and promoting phagophore-endoplasmic reticulum contacts during the lipid transfer phase of mitophagy (PubMed:33773106).

Cellular Location

Cytoplasm. Cytoplasm, cytosol. Cytoplasmic vesicle, autophagosome. Endoplasmic reticulum. Mitochondrion. Note=Mainly localizes to the cytoplasm, including cytosol (PubMed:29165041). A samll potion localizes to mitochondria; phosphorylation at Ser-34 promotes localization to mitochondria (PubMed:29165041).

APG4B Antibody (Center G254) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

APG4B Antibody (Center G254) Blocking peptide - Images**APG4B Antibody (Center G254) Blocking peptide - Background**

Autophagy is the process by which endogenous proteins and damaged organelles are destroyed intracellularly. Autophagy is postulated to be essential for cell homeostasis and cell remodeling during differentiation, metamorphosis, non-apoptotic cell death, and aging. Reduced levels of autophagy have been described in some malignant tumors, and a role for autophagy in controlling the unregulated cell growth linked to cancer has been proposed. This gene encodes a member of the autophagin protein family. The encoded protein is also designated as a member of the C-54 family of cysteine proteases. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq].

APG4B Antibody (Center G254) Blocking peptide - References

Satoo, K., et al. EMBO J. 28(9):1341-1350(2009) Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007) Ewing, R.M., et al. Mol. Syst. Biol. 3, 89 (2007) :Olsen, J.V., et al. Cell 127(3):635-648(2006) Olsen, J.V., et al. Cell 127(3):635-648(2006)