

TRAF6 Antibody(Center) Blocking peptide
Synthetic peptide
Catalog # BP19545c**Specification**

TRAF6 Antibody(Center) Blocking peptide - Product InformationPrimary Accession [Q9Y4K3](#)**TRAF6 Antibody(Center) Blocking peptide - Additional Information****Gene ID** 7189**Other Names**

TNF receptor-associated factor 6, 632-, E3 ubiquitin-protein ligase TRAF6, Interleukin-1 signal transducer, RING finger protein 85, TRAF6, RNF85

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

TRAF6 Antibody(Center) Blocking peptide - Protein Information**Name** TRAF6**Synonyms** RNF85**Function**

E3 ubiquitin ligase that, together with UBE2N and UBE2V1, mediates the synthesis of 'Lys-63'-linked-polyubiquitin chains conjugated to proteins, such as ECSIT, IKBKG, IRAK1, AKT1 and AKT2 (PubMed: [11057907](http://www.uniprot.org/citations/11057907), PubMed: [18347055](http://www.uniprot.org/citations/18347055), PubMed: [19465916](http://www.uniprot.org/citations/19465916), PubMed: [19713527](http://www.uniprot.org/citations/19713527), PubMed: [27746020](http://www.uniprot.org/citations/27746020), PubMed: [31620128](http://www.uniprot.org/citations/31620128)). Also mediates ubiquitination of free/unanchored polyubiquitin chain that leads to MAP3K7 activation (PubMed: [19675569](http://www.uniprot.org/citations/19675569)). Leads to the activation of NF-kappa-B and JUN (PubMed: [16378096](http://www.uniprot.org/citations/16378096), PubMed: [17135271](http://www.uniprot.org/citations/17135271), PubMed: [17703191](http://www.uniprot.org/citations/17703191)). Seems to also play a role in dendritic cells (DCs) maturation

and/or activation (By similarity). Represses c-Myb-mediated transactivation, in B-lymphocytes (PubMed:18093978, PubMed:18758450). Adapter protein that seems to play a role in signal transduction initiated via TNF receptor, IL-1 receptor and IL-17 receptor (PubMed:12140561, PubMed:19825828, PubMed:8837778). Regulates osteoclast differentiation by mediating the activation of adapter protein complex 1 (AP-1) and NF-kappa-B, in response to RANK-L stimulation (By similarity). Together with MAP3K8, mediates CD40 signals that activate ERK in B-cells and macrophages, and thus may play a role in the regulation of immunoglobulin production (By similarity). Acts as a regulator of the JNK and NF-kappa-B signaling pathways by initiating assembly of heterotypic 'Lys-63'-/'Lys-48'-linked branched ubiquitin chains that are then recognized by TAB2: TRAF6 catalyzes initial 'Lys-63'-linked-polyubiquitin chains that are then branched via 'Lys-48'-linked polyubiquitin by HUWE1 (PubMed:27746020). 'Lys-63'-/'Lys-48'-linked branched ubiquitin chains protect 'Lys-63'- linkages from CYLD deubiquitination (PubMed:27746020). Participates also in the TCR signaling by ubiquitinating LAT (PubMed:23514740, PubMed:25907557).

Cellular Location

Cytoplasm. Cytoplasm, cell cortex. Nucleus. Lipid droplet {ECO:0000250|UniProtKB:P70196}. Note=Found in the nuclei of some aggressive B-cell lymphoma cell lines as well as in the nuclei of both resting and activated T- and B-lymphocytes. Found in punctate nuclear body protein complexes. Ubiquitination may occur in the cytoplasm and sumoylation in the nucleus. RSAD2/viperin recruits it to the lipid droplet (By similarity).

Tissue Location

Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas

TRAF6 Antibody(Center) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

TRAF6 Antibody(Center) Blocking peptide - Images

TRAF6 Antibody(Center) Blocking peptide - Background

The protein encoded by this gene is a member of the TNFreceptor associated factor (TRAF) protein family. TRAF proteins are associated with, and mediate signal transduction from members of the TNF receptor superfamily. This protein mediates the signaling not only from the members of the TNF receptor superfamily, but also from the members of the Toll/IL-1 family. Signals from receptors such as CD40, TNFSF11/RANCE and IL-1 have been shown to be mediated by this protein. This protein also interacts with various protein kinases including IRAK1/IRAK, SRC and PKCzeta, which provides a link between distinct signaling pathways. This protein functions as a signal transducer in the NF-kappaB pathway that activates I kappa B kinase (IKK) in response to proinflammatory cytokines. The interaction of this protein with UBE2N/UBC13, and UBE2V1/UEV1A, which are ubiquitin conjugating enzymes catalyzing the formation of polyubiquitin chains, has been found to be required for IKK activation by this protein. Two alternatively spliced transcript variants encoding identical proteins have been reported. [provided by RefSeq].

TRAF6 Antibody(Center) Blocking peptide - References

Silva, L.K., et al. Eur. J. Hum. Genet. 18(11):1221-1227(2010)Hinz, M., et al. Mol. Cell
40(1):63-74(2010)Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)Bailey, S.D., et al. Diabetes
Care 33(10):2250-2253(2010)Stachon, P., et al. PLoS ONE 5 (7), E11589 (2010) :