

Mes-4 Antibody (Center) Blocking Peptide
Synthetic peptide
Catalog # BP1965c**Specification**

Mes-4 Antibody (Center) Blocking Peptide - Product InformationPrimary Accession [Q9NH52](#)**Mes-4 Antibody (Center) Blocking Peptide - Additional Information****Gene ID** 179824**Other Names**

Histone-lysine N-methyltransferase mes-4, Maternal-effect sterile protein 4, mes-4

Target/Specificity

The synthetic peptide sequence used to generate the antibody [AP1965c](/product/products/AP1965c) was selected from the Center region of human Mes-4. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Mes-4 Antibody (Center) Blocking Peptide - Protein Information**Name** mes-4 {ECO:0000312|WormBase:Y2H9A.1}**Function**

Histone methyltransferase. Dimethylates 'Lys-36' of histone H3, a specific tag for epigenetic transcriptional activation. Plays a central role in early development and is responsible for all H3 'Lys- 36' dimethylation until about the 40-cell stage. Indirectly involved in the global inactivation of the X chromosomes in germline cells, possibly by excluding the mes-2-mes-3-mes-6 repressive Polycomb complex from the autosomes (PubMed: [12077420](http://www.uniprot.org/citations/12077420), PubMed: [16968818](http://www.uniprot.org/citations/16968818)). Not related to transcription elongation (PubMed: [12077420](http://www.uniprot.org/citations/12077420), PubMed: [16968818](http://www.uniprot.org/citations/16968818)). Required for small-RNA-induced H3K27 trimethylation (PubMed: [26365259](http://www.uniprot.org/citations/26365259)). May suppress sensitivity to RNAi (PubMed: [16507136](http://www.uniprot.org/citations/16507136)).

target="_blank">16507136). May regulate the expression of genes required for vulval development (PubMed:16507136, PubMed:16710447).

Cellular Location

Nucleus. Chromosome. Note=Specifically associated with the autosomes and with the distal tip of chromosome X. Colocalizes with methylated 'Lys-4' of histone H3

Tissue Location

In adults, it is predominantly expressed in the germline, and weakly expressed in intestinal cells

Mes-4 Antibody (Center) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Mes-4 Antibody (Center) Blocking Peptide - Images**Mes-4 Antibody (Center) Blocking Peptide - Background**

Mes-4 is a histone methyltransferase. This protein dimethylates 'Lys-36' of histone H3, a specific tag for epigenetic transcriptional activation. Mes-4 plays a central role in early development and is responsible for all H3 'Lys-36' dimethylation until about the 40-cell stage. It appears to be indirectly involved in the global inactivation of the X chromosomes in germ line cells, possibly by excluding the mes-2-mes-3-mes-6 repressive Polycomb complex from the autosomes.