

## JUN Antibody (C-term E248) Blocking peptide

Synthetic peptide Catalog # BP1984k

## **Specification**

## JUN Antibody (C-term E248) Blocking peptide - Product Information

Primary Accession P05412
Other Accession NP 002219

## JUN Antibody (C-term E248) Blocking peptide - Additional Information

### **Gene ID 3725**

### **Other Names**

Transcription factor AP-1, Activator protein 1, AP1, Proto-oncogene c-Jun, V-jun avian sarcoma virus 17 oncogene homolog, p39, JUN

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

# JUN Antibody (C-term E248) Blocking peptide - Protein Information

## Name JUN

### **Function**

Transcription factor that recognizes and binds to the AP-1 consensus motif 5'-TGA[GC]TCA-3' (PubMed:<a href="http://www.uniprot.org/citations/10995748" target=" blank">10995748</a>, PubMed: <a href="http://www.uniprot.org/citations/22083952" target="blank">22083952</a>). Heterodimerizes with proteins of the FOS family to form an AP-1 transcription complex, thereby enhancing its DNA binding activity to the AP-1 consensus sequence 5'-TGA[GC]TCA-3' and enhancing its transcriptional activity (By similarity). Together with FOSB, plays a role in activation-induced cell death of T cells by binding to the AP-1 promoter site of FASLG/CD95L, and inducing its transcription in response to activation of the TCR/CD3 signaling pathway (PubMed:<a href="http://www.uniprot.org/citations/12618758" target=" blank">12618758</a>). Promotes activity of NR5A1 when phosphorylated by HIPK3 leading to increased steroidogenic gene expression upon cAMP signaling pathway stimulation (PubMed:<a href="http://www.uniprot.org/citations/17210646" target=" blank">17210646</a>). Involved in activated KRAS-mediated transcriptional activation of USP28 in colorectal cancer (CRC) cells (PubMed:<a href="http://www.uniprot.org/citations/24623306" target=" blank">24623306</a>). Binds to the USP28 promoter in colorectal cancer (CRC) cells (PubMed:<a href="http://www.uniprot.org/citations/24623306" target=" blank">24623306</a>).



**Cellular Location** Nucleus.

### **Tissue Location**

Expressed in the developing and adult prostate and prostate cancer cells.

## JUN Antibody (C-term E248) Blocking peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

### • Blocking Peptides

JUN Antibody (C-term E248) Blocking peptide - Images

# JUN Antibody (C-term E248) Blocking peptide - Background

JUN is the putative transforming gene of aviansarcoma virus 17. It encodes a protein which is highly similar tothe viral protein, and which interacts directly with specifictarget DNA sequences to regulate gene expression. This gene isintronless and is mapped to 1p32-p31, a chromosomal region involved in both translocations and deletions in human malignancies.

## JUN Antibody (C-term E248) Blocking peptide - References

Lin, A., et al. Cell 70(5):777-789(1992)Lin, A., et al. Cell 70(5):777-789(1992)Chevray, P.M., et al. Proc. Natl. Acad. Sci. U.S.A. 89(13):5789-5793(1992)Bengal, E., et al. Cell 68(3):507-519(1992)Alvarez, E., et al. J. Biol. Chem. 266(23):15277-15285(1991)