

SELM Blocking Peptide (C-term)
Synthetic peptide
Catalog # BP20147b**Specification**

SELM Blocking Peptide (C-term) - Product Information

Primary Accession [Q8WWX9](#)
Other Accession [NP_536355.1](#)

SELM Blocking Peptide (C-term) - Additional Information

Gene ID 140606

Other Names

Selenoprotein M, SelM, SELM, SEPM

Target/Specificity

The synthetic peptide sequence is selected from aa 132-145 of HUMAN SELM

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

SELM Blocking Peptide (C-term) - Protein Information

Name SELENOM {ECO:0000303|PubMed:27645994, ECO:0000312|HGNC:HGNC:30397}

Function

May function as a thiol-disulfide oxidoreductase that participates in disulfide bond formation.

Cellular Location

Cytoplasm, perinuclear region. Endoplasmic reticulum. Golgi apparatus. Note=Localized to perinuclear structures corresponding to Golgi and endoplasmic reticulum

Tissue Location

Widely expressed..

SELM Blocking Peptide (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

SELM Blocking Peptide (C-term) - Images

SELM Blocking Peptide (C-term) - Background

This gene encodes a selenoprotein, which contains a selenocysteine (Sec) residue at its active site. The selenocysteine is encoded by the UGA codon that normally signals translation termination. The 3' UTR of selenoprotein genes have a common stem-loop structure, the sec insertion sequence (SECIS), that is necessary for the recognition of UGA as a Sec codon rather than as a stop signal. This gene is expressed in a variety of tissues, and the protein is localized to the perinuclear structures. [provided by RefSeq].

SELM Blocking Peptide (C-term) - References

Korotkov, K.V., et al. Mol. Cell. Biol. 22(5):1402-1411(2002)