

**PPP2CA Blocking Peptide (Center)**  
**Synthetic peptide**  
**Catalog # BP20181c****Specification**

---

**PPP2CA Blocking Peptide (Center) - Product Information**

Primary Accession [P67775](#)  
Other Accession [P23696](#), [P63331](#), [P67777](#), [P67776](#), [P63330](#),  
[P67774](#), [NP\\_002706.1](#)

**PPP2CA Blocking Peptide (Center) - Additional Information**

**Gene ID** 5515

**Other Names**

Serine/threonine-protein phosphatase 2A catalytic subunit alpha isoform, PP2A-alpha, Replication protein C, RP-C, PPP2CA

**Target/Specificity**

The synthetic peptide sequence is selected from aa 103-116 of HUMAN PPP2CA

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**PPP2CA Blocking Peptide (Center) - Protein Information**

**Name** PPP2CA

**Function**

PP2A is the major phosphatase for microtubule-associated proteins (MAPs) (PubMed:<a href="http://www.uniprot.org/citations/22613722" target="\_blank">22613722</a>). PP2A can modulate the activity of phosphorylase B kinase casein kinase 2, mitogen-stimulated S6 kinase, and MAP-2 kinase (PubMed:<a href="http://www.uniprot.org/citations/22613722" target="\_blank">22613722</a>). Cooperates with SGO2 to protect centromeric cohesin from separase-mediated cleavage in oocytes specifically during meiosis I (By similarity). Can dephosphorylate SV40 large T antigen and p53/TP53 (PubMed:<a href="http://www.uniprot.org/citations/17245430" target="\_blank">17245430</a>). Activates RAF1 by dephosphorylating it at 'Ser-259' (PubMed:<a href="http://www.uniprot.org/citations/10801873" target="\_blank">10801873</a>). Mediates dephosphorylation of WEE1, preventing its ubiquitin-mediated proteolysis, increasing WEE1 protein levels, and promoting the G2/M checkpoint (PubMed:<a

[33108758](http://www.uniprot.org/citations/33108758)). Mediates dephosphorylation of MYC; promoting its ubiquitin-mediated proteolysis: interaction with AMBRA1 enhances interaction between PPP2CA and MYC (PubMed:<[25438055](http://www.uniprot.org/citations/25438055)>). Mediates dephosphorylation of FOXO3; promoting its stabilization: interaction with AMBRA1 enhances interaction between PPP2CA and FOXO3 (PubMed:<[30513302](http://www.uniprot.org/citations/30513302)>). Catalyzes dephosphorylation of the pyrin domain of NLRP3, promoting assembly of the NLRP3 inflammasome (By similarity).

#### **Cellular Location**

Cytoplasm. Nucleus. Chromosome, centromere. Cytoplasm, cytoskeleton, spindle pole. Note=In prometaphase cells, but not in anaphase cells, localizes at centromeres (PubMed:16541025). During mitosis, also found at spindle poles (PubMed:16541025). Centromeric localization requires the presence of SGO2 (By similarity) {ECO:0000250|UniProtKB:P63330, ECO:0000269|PubMed:16541025}

#### **PPP2CA Blocking Peptide (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

#### **PPP2CA Blocking Peptide (Center) - Images**

#### **PPP2CA Blocking Peptide (Center) - Background**

This gene encodes the phosphatase 2A catalytic subunit. Protein phosphatase 2A is one of the four major Ser/Thr phosphatases, and it is implicated in the negative control of cell growth and division. It consists of a common heteromeric core enzyme, which is composed of a catalytic subunit and a constant regulatory subunit, that associates with a variety of regulatory subunits. This gene encodes an alpha isoform of the catalytic subunit.

#### **PPP2CA Blocking Peptide (Center) - References**

Shimada, M., et al. Hum. Genet. 128(4):433-441(2010)  
Jayadeva, G., et al. J. Biol. Chem. 285(39):29863-29873(2010)  
Pradhan, S., et al. J. Biol. Chem. 285(38):29059-29068(2010)  
Schmitz, M.H., et al. Nat. Cell Biol. 12(9):886-893(2010)  
Antony, R., et al. J. Biol. Chem. 285(24):18301-18308(2010)