

DENR Blocking Peptide (Center)

Synthetic peptide Catalog # BP20311c

Specification

DENR Blocking Peptide (Center) - Product Information

Primary Accession <u>043583</u>

Other Accession Q9CQI6, Q2HJ47

DENR Blocking Peptide (Center) - Additional Information

Gene ID 8562

Other Names

Density-regulated protein, DRP, Protein DRP1, Smooth muscle cell-associated protein 3, SMAP-3, DENR, DRP1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

DENR Blocking Peptide (Center) - Protein Information

Name DENR

Synonyms DRP1


Function

May be involved in the translation of target mRNAs by scanning and recognition of the initiation codon. Involved in translation initiation; promotes recruitmnet of aminoacetyled initiator tRNA to P site of 40S ribosomes. Can promote release of deacylated tRNA and mRNA from recycled 40S subunits following ABCE1-mediated dissociation of post-termination ribosomal complexes into subunits. Plays a role in the modulation of the translational profile of a subset of cancer-related mRNAs when recruited to the translational initiation complex by the oncogene MCTS1.

Tissue Location

Highly expressed in heart and skeletal muscle and moderately expressed in the brain, placenta, liver and pancreas. Weakly expressed in the lung and kidney.

DENR Blocking Peptide (Center) - Protocols





Tel: 858.875.1900 Fax: 858.875.1999

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

DENR Blocking Peptide (Center) - Images

DENR Blocking Peptide (Center) - Background

May be involved in the translation of target mRNAs by scanning and recognition of the initiation codon. Plays a role in the modulation of the translational profile of a subset of cancer-related mRNAs when recruited to the translational initiation complex by the oncogene MCTS1.