

REG3A Blocking Peptide (C-term)

Synthetic peptide

Catalog # BP20597c

Specification

REG3A Blocking Peptide (C-term) - Product Information

Primary Accession

[Q06141](#)**REG3A Blocking Peptide (C-term) - Additional Information**

Gene ID 5068

Other Names

Regenerating islet-derived protein 3-alpha, REG-3-alpha, Hepatointestinal pancreatic protein, HIP/PAP, Human proislet peptide, Pancreatitis-associated protein 1, Regenerating islet-derived protein III-alpha, Reg III-alpha, Regenerating islet-derived protein 3-alpha 165 kDa form, Regenerating islet-derived protein 3-alpha 15 kDa form, REG3A, HIP, PAP, PAP1

Target/Specificity

The synthetic peptide sequence is selected from aa 149-163 of HUMAN REG3A

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

REG3A Blocking Peptide (C-term) - Protein InformationName REG3A ([HGNC:8601](#))**Function**

[Regenerating islet-derived protein 3-alpha 15 kDa form]: Bactericidal C-type lectin which acts exclusively against Gram-positive bacteria and mediates bacterial killing by binding to surface-exposed carbohydrate moieties of peptidoglycan (PubMed:16931762). Binds membrane phospholipids and kills bacteria by forming a hexameric membrane-permeabilizing oligomeric pore (PubMed:24256734).

Cellular Location

Secreted. Note=Found in the apical region of pancreatic acinar cells

Tissue Location

Expressed by keratinocytes (PubMed:27830702). Highly expressed in epidermal keratinocytes of psoriasis patients (at protein level) (PubMed:22727489). Constitutively expressed in intestine. Low expression is found in healthy pancreas. Overexpressed during the acute phase of pancreatitis and in some patients with chronic pancreatitis (PubMed:1469087).

REG3A Blocking Peptide (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

REG3A Blocking Peptide (C-term) - Images

REG3A Blocking Peptide (C-term) - Background

Bactericidal C-type lectin which acts exclusively against Gram-positive bacteria and mediates bacterial killing by binding to surface-exposed carbohydrate moieties of peptidoglycan. Regulates keratinocyte proliferation and differentiation after skin injury via activation of EXTL3-PI3K-AKT signaling pathway.

REG3A Blocking Peptide (C-term) - References

Itoh T.,et al.Biochim. Biophys. Acta 1172:184-186(1993).
Orelle B.,et al.J. Clin. Invest. 90:2284-2291(1992).
Lasserre C.,et al.Cancer Res. 52:5089-5095(1992).
Duseti N.J.,et al.Genomics 19:108-114(1994).
Lasserre C.,et al.Eur. J. Biochem. 224:29-38(1994).