

Histone H2A.X (Ser139) Blocking Peptide
Synthetic peptide
Catalog # BP20703b**Specification**

Histone H2A.X (Ser139) Blocking Peptide - Product InformationPrimary Accession [P16104](#)**Histone H2A.X (Ser139) Blocking Peptide - Additional Information****Gene ID** 3014**Other Names**

Histone H2AX, H2a/x, Histone H2AX, H2AFX, H2AX

Target/Specificity

The synthetic peptide sequence is selected from aa 134-143 of HUMAN H2AFX

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

Histone H2A.X (Ser139) Blocking Peptide - Protein Information**Name** H2AX ([HGNC:4739](#))**Function**

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post- translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically when modified by C-terminal phosphorylation.

Cellular Location

Nucleus. Chromosome

Histone H2A.X (Ser139) Blocking Peptide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

Histone H2A.X (Ser139) Blocking Peptide - Images

Histone H2A.X (Ser139) Blocking Peptide - Background

Variant histone H2A which replaces conventional H2A in a subset of nucleosomes. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling. Required for checkpoint-mediated arrest of cell cycle progression in response to low doses of ionizing radiation and for efficient repair of DNA double strand breaks (DSBs) specifically when modified by C- terminal phosphorylation.

Histone H2A.X (Ser139) Blocking Peptide - References

Mannironi C.,et al.Nucleic Acids Res. 17:9113-9126(1989).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Rogakou E.P.,et al.J. Biol. Chem. 273:5858-5868(1998).
Rogakou E.P.,et al.J. Cell Biol. 146:905-916(1999).
Paull T.T.,et al.Curr. Biol. 10:886-895(2000).