

## PRPF6 Blocking Peptide (N-term)

Synthetic peptide Catalog # BP20777a

### **Specification**

## PRPF6 Blocking Peptide (N-term) - Product Information

Primary Accession <u>094906</u>

Other Accession A1A5S1, Q91YR7, Q2KIIO

# PRPF6 Blocking Peptide (N-term) - Additional Information

**Gene ID 24148** 

#### **Other Names**

Pre-mRNA-processing factor 6, Androgen receptor N-terminal domain-transactivating protein 1, ANT-1, PRP6 homolog, U5 snRNP-associated 102 kDa protein, U5-102 kDa protein, PRPF6, C20orf14

#### Target/Specificity

The synthetic peptide sequence is selected from aa 196-209 of HUMAN PRPF6

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## PRPF6 Blocking Peptide (N-term) - Protein Information

Name PRPF6

Synonyms C20orf14

### **Function**

Involved in pre-mRNA splicing as component of the U4/U6-U5 tri-snRNP complex, one of the building blocks of the spliceosome (PubMed:<a href="http://www.uniprot.org/citations/28781166" target="\_blank">28781166</a>, PubMed:<a href="http://www.uniprot.org/citations/21549338" target="\_blank">21549338</a>). Enhances dihydrotestosterone- induced transactivation activity of AR, as well as dexamethasone- induced transactivation activity of NR3C1, but does not affect estrogen-induced transactivation.

### **Cellular Location**

Nucleus, nucleoplasm. Nucleus speckle. Note=Localized in splicing speckles.



**Tissue Location** Widely expressed...

## PRPF6 Blocking Peptide (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

PRPF6 Blocking Peptide (N-term) - Images

PRPF6 Blocking Peptide (N-term) - Background

Involved in pre-mRNA splicing as component of the U4/U6- U5 tri-snRNP complex, one of the building blocks of the spliceosome. Enhances dihydrotestosterone-induced transactivation activity of AR, as well as dexamethasone-induced transactivation activity of NR3C1, but does not affect estrogen-induced transactivation.

### PRPF6 Blocking Peptide (N-term) - References

Nishikimi A.,et al.Biochim. Biophys. Acta 1435:147-152(1999). Makarov E.M.,et al.J. Mol. Biol. 298:567-575(2000). Yuryev A.,et al.Mol. Cell. Biol. 20:4870-4878(2000). Ota T.,et al.Nat. Genet. 36:40-45(2004). Deloukas P.,et al.Nature 414:865-871(2001).