

**Mouse Csk Blocking Peptide (N-term)**

Synthetic peptide

Catalog # BP20789a

**Specification**

---

**Mouse Csk Blocking Peptide (N-term) - Product Information**

Primary Accession

[P41241](#)

Other Accession

[P32577](#), [P41240](#), [Q0VBZ0](#)**Mouse Csk Blocking Peptide (N-term) - Additional Information****Gene ID** 12988**Other Names**

Tyrosine-protein kinase CSK, C-Src kinase, Protein-tyrosine kinase MPK-2, p50CSK, Csk

**Target/Specificity**

The synthetic peptide sequence is selected from aa 67-81 of HUMAN Csk

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**Mouse Csk Blocking Peptide (N-term) - Protein Information****Name** Csk**Function**

Non-receptor tyrosine-protein kinase that plays an important role in the regulation of cell growth, differentiation, migration and immune response. Phosphorylates tyrosine residues located in the C-terminal tails of Src-family kinases (SFKs) including LCK, SRC, HCK, FYN, LYN, CSK or YES1. Upon tail phosphorylation, Src-family members engage in intramolecular interactions between the phosphotyrosine tail and the SH2 domain that result in an inactive conformation. To inhibit SFKs, CSK is recruited to the plasma membrane via binding to transmembrane proteins or adapter proteins located near the plasma membrane. Suppresses signaling by various surface receptors, including T-cell receptor (TCR) and B-cell receptor (BCR) by phosphorylating and maintaining inactive several positive effectors such as FYN or LCK (By similarity).

**Cellular Location**

Cytoplasm. Cell membrane. Note=Mainly cytoplasmic, also present in lipid rafts

**Tissue Location**

Ubiquitous, but most abundant in thymus and spleen, as well as in neonatal brain

### **Mouse Csk Blocking Peptide (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **Mouse Csk Blocking Peptide (N-term) - Images**

### **Mouse Csk Blocking Peptide (N-term) - Background**

Non-receptor tyrosine-protein kinase that plays an important role in the regulation of cell growth, differentiation, migration and immune response. Phosphorylates tyrosine residues located in the C-terminal tails of Src-family kinases (SFKs) including LCK, SRC, HCK, FYN, LYN or YES1. Upon tail phosphorylation, Src-family members engage in intramolecular interactions between the phosphotyrosine tail and the SH2 domain that result in an inactive conformation. To inhibit SFKs, CSK is recruited to the plasma membrane via binding to transmembrane proteins or adapter proteins located near the plasma membrane. Suppresses signaling by various surface receptors, including T- cell receptor (TCR) and B-cell receptor (BCR) by phosphorylating and maintaining inactive several positive effectors such as FYN or LCK (By similarity).

### **Mouse Csk Blocking Peptide (N-term) - References**

Klages S.,et al.Proc. Natl. Acad. Sci. U.S.A. 91:2597-2601(1994).  
Farber C.R.,et al.Submitted (JAN-2005) to the EMBL/GenBank/DDBJ databases.  
Carninci P.,et al.Science 309:1559-1563(2005).  
Gilardi-Hebenstreit P.,et al.Oncogene 7:2499-2506(1992).  
Lubec G.,et al.Submitted (JAN-2009) to UniProtKB.