

SMAD1 Blocking Peptide (Center)
Synthetic peptide
Catalog # BP20916a**Specification**

SMAD1 Blocking Peptide (Center) - Product Information

Primary Accession [Q15797](#)
Other Accession [P97588](#), [P70340](#), [Q1JQA2](#)

SMAD1 Blocking Peptide (Center) - Additional Information

Gene ID 4086

Other Names

Mothers against decapentaplegic homolog 1, MAD homolog 1, Mothers against DPP homolog 1, JV4-1, Mad-related protein 1, SMAD family member 1, SMAD 1, Smad1, hSMAD1, Transforming growth factor-beta-signaling protein 1, BSP-1, SMAD1, BSP1, MADH1, MADR1

Target/Specificity

The synthetic peptide sequence is selected from aa 257-270 of HUMAN SMAD1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

SMAD1 Blocking Peptide (Center) - Protein Information

Name SMAD1

Synonyms BSP1, MADH1, MADR1

Function

Transcriptional modulator that plays a role in various cellular processes, including embryonic development, cell differentiation, and tissue homeostasis (PubMed:[9335504](http://www.uniprot.org/citations/9335504)). Upon BMP ligand binding to their receptors at the cell surface, is phosphorylated by activated type I BMP receptors (BMPRI) and associates with SMAD4 to form an heteromeric complex which translocates into the nucleus acting as transcription factor (PubMed:[33667543](http://www.uniprot.org/citations/33667543)). In turn, the hetero-trimeric complex recognizes cis-regulatory elements containing Smad Binding Elements (SBEs) to modulate the outcome of the signaling network (PubMed:[33667543](http://www.uniprot.org/citations/33667543)).

SMAD1/OAZ1/PSMB4 complex mediates the degradation of the CREBBP/EP300 repressor SNIP1. Positively regulates BMP4-induced expression of odontogenic development regulator MSX1 following IPO7-mediated nuclear import (By similarity).

Cellular Location

Cytoplasm. Nucleus Note=Cytoplasmic in the absence of ligand. Migrates to the nucleus when complexed with SMAD4 (PubMed:15647271). Co-localizes with LEMD3 at the nucleus inner membrane (PubMed:15647271). Exported from the nucleus to the cytoplasm when dephosphorylated (By similarity) {ECO:0000250|UniProtKB:P70340, ECO:0000269|PubMed:15647271}

Tissue Location

Ubiquitous. Highest expression seen in the heart and skeletal muscle

SMAD1 Blocking Peptide (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

SMAD1 Blocking Peptide (Center) - Images**SMAD1 Blocking Peptide (Center) - Background**

Transcriptional modulator activated by BMP (bone morphogenetic proteins) type 1 receptor kinase. SMAD1 is a receptor-regulated SMAD (R-SMAD). SMAD1/OAZ1/PSMB4 complex mediates the degradation of the CREBBP/EP300 repressor SNIP1.

SMAD1 Blocking Peptide (Center) - References

Riggins G.J.,et al.Nat. Genet. 13:347-349(1996).
Liu F.,et al.Nature 381:620-623(1996).
Hoodless P.A.,et al.Cell 85:489-500(1996).
Lechleider R.J.,et al.J. Biol. Chem. 271:17617-17620(1996).
Zhang Y.,et al.Nature 383:168-172(1996).