

PDX1 Blocking Peptide (N-term)
Synthetic peptide
Catalog # BP20965b**Specification**

PDX1 Blocking Peptide (N-term) - Product InformationPrimary Accession [P52945](#)**PDX1 Blocking Peptide (N-term) - Additional Information****Gene ID** 3651**Other Names**

Pancreas/duodenum homeobox protein 1, PDX-1, Glucose-sensitive factor, GSF, Insulin promoter factor 1, IPF-1, Insulin upstream factor 1, IUF-1, Islet/duodenum homeobox-1, IDX-1, Somatostatin-transactivating factor 1, STF-1, PDX1, IPF1, STF1

Target/Specificity

The synthetic peptide sequence is selected from aa 21-35 of HUMAN PDX1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

PDX1 Blocking Peptide (N-term) - Protein Information**Name** PDX1**Synonyms** IPF1, STF1**Function**

Activates insulin, somatostatin, glucokinase, islet amyloid polypeptide and glucose transporter type 2 gene transcription. Particularly involved in glucose-dependent regulation of insulin gene transcription. As part of a PDX1:PBX1b:MEIS2b complex in pancreatic acinar cells is involved in the transcriptional activation of the ELA1 enhancer; the complex binds to the enhancer B element and cooperates with the transcription factor 1 complex (PTF1) bound to the enhancer A element. Binds preferentially the DNA motif 5'-[CT]TAAT[TG]-3'. During development, specifies the early pancreatic epithelium, permitting its proliferation, branching and subsequent differentiation. At adult stage, required for maintaining the hormone-producing phenotype of the beta-cell.

Cellular Location

Nucleus. Cytoplasm, cytosol.

Tissue Location

Duodenum and pancreas (Langerhans islet beta cells and small subsets of endocrine non-beta-cells, at low levels in acinar cells)

PDX1 Blocking Peptide (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

PDX1 Blocking Peptide (N-term) - Images**PDX1 Blocking Peptide (N-term) - Background**

Activates insulin, somatostatin, glucokinase, islet amyloid polypeptide and glucose transporter type 2 gene transcription. Particularly involved in glucose-dependent regulation of insulin gene transcription. As part of a PDX1:PBX1b:MEIS2b complex in pancreatic acinar cells is involved in the transcriptional activation of the ELA1 enhancer; the complex binds to the enhancer B element and cooperates with the transcription factor 1 complex (PTF1) bound to the enhancer A element. Binds preferentially the DNA motif 5'-[CT]TAAT[TG]-3'. During development, specifies the early pancreatic epithelium, permitting its proliferation, branching and subsequent differentiation. At adult stage, required for maintaining the hormone-producing phenotype of the beta-cell.

PDX1 Blocking Peptide (N-term) - References

Stoffel M.,et al.Genomics 28:125-126(1995).
Inoue H.,et al.Diabetes 45:789-794(1996).
Hiroshi I.,et al.Submitted (JUN-1995) to the EMBL/GenBank/DDBJ databases.
Marshak S.,et al.Submitted (AUG-1996) to the EMBL/GenBank/DDBJ databases.
Hara M.,et al.Submitted (DEC-1997) to the EMBL/GenBank/DDBJ databases.