

CHST11 Blocking Peptide (Center)

Synthetic peptide

Catalog # BP21059a

Specification

CHST11 Blocking Peptide (Center) - Product Information

Primary Accession

[O9NPF2](#)

Other Accession

[P69478](#), [Q9JME2](#)**CHST11 Blocking Peptide (Center) - Additional Information**

Gene ID 50515

Other Names

Carbohydrate sulfotransferase 11, Chondroitin 4-O-sulfotransferase 1, Chondroitin 4-sulfotransferase 1, C4S-1, C4ST-1, C4ST1, CHST11

Target/Specificity

The synthetic peptide sequence is selected from aa 246-260 of HUMAN CHST11

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CHST11 Blocking Peptide (Center) - Protein Information

Name CHST11

Function

Catalyzes the transfer of sulfate to position 4 of the N- acetylgalactosamine (GalNAc) residue of chondroitin. Chondroitin sulfate constitutes the predominant proteoglycan present in cartilage and is distributed on the surfaces of many cells and extracellular matrices. Can also sulfate Gal residues in desulfated dermatan sulfate. Preferentially sulfates in GlcA->GalNAc unit than in IdoA->GalNAc unit. Does not form 4, 6-di-O-sulfated GalNAc when chondroitin sulfate C is used as an acceptor.

Cellular Location

Golgi apparatus membrane; Single-pass type II membrane protein

Tissue Location

Widely expressed. Highly expressed in spleen, thymus, bone marrow, peripheral blood leukocytes, lymph node, heart, brain, lung and placenta.

CHST11 Blocking Peptide (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

CHST11 Blocking Peptide (Center) - Images

CHST11 Blocking Peptide (Center) - Background

Catalyzes the transfer of sulfate to position 4 of the N-acetylgalactosamine (GalNAc) residue of chondroitin. Chondroitin sulfate constitutes the predominant proteoglycan present in cartilage and is distributed on the surfaces of many cells and extracellular matrices. Can also sulfate Gal residues in desulfated dermatan sulfate. Preferentially sulfates in GlcA->GalNAc unit than in IdoA->GalNAc unit. Does not form 4, 6- di-O-sulfated GalNAc when chondroitin sulfate C is used as an acceptor.

CHST11 Blocking Peptide (Center) - References

Okuda T.,et al.J. Biochem. 128:763-770(2000).
Hiraoka N.,et al.J. Biol. Chem. 275:20188-20196(2000).
Xia G.,et al.Submitted (MAY-2000) to the EMBL/GenBank/DDBJ databases.
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Mural R.J.,et al.Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.