

**UBCE7IP5 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP2109b****Specification**

---

**UBCE7IP5 Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [O94941](#)**UBCE7IP5 Antibody (C-term) Blocking Peptide - Additional Information****Gene ID** 22888**Other Names**

RING finger protein 37, 632-, UBOX5 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=17777](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=17777))  
HGNC:17777

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP2109b](/product/products/AP2109b) was selected from the C-term region of human UBCE7IP5. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**UBCE7IP5 Antibody (C-term) Blocking Peptide - Protein Information****Name** UBOX5 ([HGNC:17777](#))**Function**

May have a ubiquitin-protein ligase activity acting as an E3 ubiquitin-protein ligase or as a ubiquitin-ubiquitin ligase promoting elongation of ubiquitin chains on substrates.

**Cellular Location**

Nucleus. Note=Enriched in nuclear bodies

**Tissue Location**

Expressed in liver, heart, brain, kidney and testis.

## **UBCE7IP5 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

## **UBCE7IP5 Antibody (C-term) Blocking Peptide - Images**

## **UBCE7IP5 Antibody (C-term) Blocking Peptide - Background**

Ubiquitin is a 76 amino acid highly conserved eukaryotic polypeptide that selectively marks cellular proteins for proteolytic degradation by the 26S proteasome. The process of target selection, covalent attachment and shuttle to the 26S proteasome is a vital means of regulating the concentrations of key regulatory proteins in the cell by limiting their lifespans. Polyubiquitination is a common feature of this modification. Serial steps for modification include the activation of ubiquitin, an ATP-dependent formation of a thioester bond between ubiquitin and the enzyme E1, transfer by transacylation of ubiquitin from E1 to the ubiquitin conjugating enzyme E2, and covalent linkage to the target protein directly by E2 or via E3 ligase enzyme. Deubiquitination enzymes also exist to reverse the marking of protein substrates. Posttranslational tagging by Ub is involved in a multitude of cellular processes, including the cell cycle, cell growth and differentiation, embryogenesis, apoptosis, signal transduction, DNA repair, regulation of transcription and DNA replication, transmembrane transport, stress responses, the immune response, and nervous system functions.

## **UBCE7IP5 Antibody (C-term) Blocking Peptide - References**

Martinez-Noel, G., et al., FEBS Lett. 454(3):257-261 (1999).