

**UBE4B Antibody (C-term) Blocking Peptide**  
Synthetic peptide  
Catalog # BP2111b**Specification**

---

**UBE4B Antibody (C-term) Blocking Peptide - Product Information**Primary Accession [O95155](#)**UBE4B Antibody (C-term) Blocking Peptide - Additional Information**

Gene ID 10277

**Other Names**

Ubiquitin conjugation factor E4 B, 632-, UBE4B (&lt;a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=12500" target="\_blank"&gt;HGNC:12500&lt;/a&gt;)

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody &lt;a href="/product/products/AP2111b"&gt;AP2111b&lt;/a&gt; was selected from the C-term region of human UBE4B . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**UBE4B Antibody (C-term) Blocking Peptide - Protein Information**Name UBE4B ([HGNC:12500](#))**Function**

Ubiquitin-protein ligase that probably functions as an E3 ligase in conjunction with specific E1 and E2 ligases (By similarity). May also function as an E4 ligase mediating the assembly of polyubiquitin chains on substrates ubiquitinated by another E3 ubiquitin ligase (By similarity). May regulate myosin assembly in striated muscles together with STUB1 and VCP/p97 by targeting myosin chaperone UNC45B for proteasomal degradation (PubMed:&lt;a href="http://www.uniprot.org/citations/17369820" target="\_blank"&gt;17369820&lt;/a&gt;).

**Cellular Location**

Cytoplasm. Nucleus {ECO:0000250|UniProtKB:Q9ES00}

**Tissue Location**

Expressed in differentiated myotubes (at protein level) (PubMed:17369820). Highest expression in ovary, testis, heart and skeletal muscle (PubMed:11802788). Expression is low in colon, thymus and peripheral blood leukocytes (PubMed:11802788). Almost undetectable in lung and spleen (PubMed:11802788)

**UBE4B Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

**UBE4B Antibody (C-term) Blocking Peptide - Images****UBE4B Antibody (C-term) Blocking Peptide - Background**

Ubiquitin is a 76 amino acid highly conserved eukaryotic polypeptide that selectively marks cellular proteins for proteolytic degradation by the 26S proteasome. The process of target selection, covalent attachment and shuttle to the 26S proteasome is a vital means of regulating the concentrations of key regulatory proteins in the cell by limiting their lifespans. Polyubiquitination is a common feature of this modification. Serial steps for modification include the activation of ubiquitin, an ATP-dependent formation of a thioester bond between ubiquitin and the enzyme E1, transfer by transacylation of ubiquitin from E1 to the ubiquitin conjugating enzyme E2, and covalent linkage to the target protein directly by E2 or via E3 ligase enzyme. Deubiquitination enzymes also exist to reverse the marking of protein substrates. Posttranslational tagging by Ub is involved in a multitude of cellular processes, including the cell cycle, cell growth and differentiation, embryogenesis, apoptosis, signal transduction, DNA repair, regulation of transcription and DNA replication, transmembrane transport, stress responses, the immune response, and nervous system functions.

**UBE4B Antibody (C-term) Blocking Peptide - References**

Ishikawa, K., et al., DNA Res. 5(3):169-176 (1998). Mahoney, J.A., et al., Biochem. J. 361 (Pt 3), 587-595 (2002).