

(DANRE) cisd2 Blocking Peptide (C-Term)
Synthetic peptide
Catalog # BP21176a

Specification

(DANRE) cisd2 Blocking Peptide (C-Term) - Product Information

Primary Accession [Q7T326](#)

(DANRE) cisd2 Blocking Peptide (C-Term) - Additional Information

Gene ID 393354

Other Names

CDGSH iron-sulfur domain-containing protein 2, cisd2

Target/Specificity

The synthetic peptide sequence is selected from aa 110-124 of HUMAN cisd2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

(DANRE) cisd2 Blocking Peptide (C-Term) - Protein Information

Name cisd2

Function

Regulator of autophagy that contributes to antagonize becn1- mediated cellular autophagy at the endoplasmic reticulum. Participates in the interaction of bcl2 with becn1 and is required for bcl2-mediated depression of endoplasmic reticulum Ca(2+) stores during autophagy (By similarity).

Cellular Location

Endoplasmic reticulum membrane; Single-pass membrane protein. Mitochondrion outer membrane; Single-pass membrane protein

(DANRE) cisd2 Blocking Peptide (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

(DANRE) cisd2 Blocking Peptide (C-Term) - Images**(DANRE) cisd2 Blocking Peptide (C-Term) - Background**

Regulator of autophagy that contributes to antagonize becn1-mediated cellular autophagy at the endoplasmic reticulum. Participates in the interaction of bcl2 with becn1 and is required for bcl2-mediated depression of endoplasmic reticulum Ca(2+) stores during autophagy (By similarity).

(DANRE) cisd2 Blocking Peptide (C-Term) - References

Song H.-D., et al. Proc. Natl. Acad. Sci. U.S.A. 101:16240-16245(2004).
Howe K., et al. Nature 496:498-503(2013).