

(Mouse) Uhrf1 Blocking Peptide (C-term)
Synthetic peptide
Catalog # BP21204b

Specification

(Mouse) Uhrf1 Blocking Peptide (C-term) - Product Information

Primary Accession [Q8VDF2](#)

(Mouse) Uhrf1 Blocking Peptide (C-term) - Additional Information

Gene ID 18140

Other Names

E3 ubiquitin-protein ligase UHRF1, 632-, Nuclear protein 95, Nuclear zinc finger protein Np95, Ubiquitin-like PHD and RING finger domain-containing protein 1, mUhrf1, Ubiquitin-like-containing PHD and RING finger domains protein 1, Uhrf1, Np95

Target/Specificity

The synthetic peptide sequence is selected from aa 596-608 of HUMAN Uhrf1

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

(Mouse) Uhrf1 Blocking Peptide (C-term) - Protein Information

Name Uhrf1

Synonyms Np95

Function

Multidomain protein that acts as a key epigenetic regulator by bridging DNA methylation and chromatin modification. Specifically recognizes and binds hemimethylated DNA at replication forks via its YDG domain and recruits DNMT1 methyltransferase to ensure faithful propagation of the DNA methylation patterns through DNA replication. In addition to its role in maintenance of DNA methylation, also plays a key role in chromatin modification: through its tudor-like regions and PHD-type zinc fingers, specifically recognizes and binds histone H3 trimethylated at 'Lys-9' (H3K9me3) and unmethylated at 'Arg-2' (H3R2me0), respectively, and recruits chromatin proteins. Enriched in pericentric heterochromatin where it recruits different chromatin modifiers required for this chromatin replication. Also localizes to euchromatic regions where it negatively regulates transcription possibly by impacting DNA methylation and histone modifications. Has E3 ubiquitin-protein ligase activity by mediating the ubiquitination of target proteins such as histone

H3 and PML. It is still unclear how E3 ubiquitin-protein ligase activity is related to its role in chromatin in vivo. Plays a role in DNA repair by cooperating with UHRF2 to ensure recruitment of FANCD2 to interstrand cross-links (ICLs) leading to FANCD2 activation. Plays a pivotal role in the establishment of correct spindle architecture by catalyzing the 'Lys-63'-linked ubiquitination of KIF11, thereby controlling KIF11 localization on the spindle.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00358, ECO:0000269|PubMed:10984098, ECO:0000269|PubMed:11161719, ECO:0000269|PubMed:14993289, ECO:0000269|PubMed:17994007, ECO:0000269|PubMed:21489993, ECO:0000269|PubMed:36056023, ECO:0000269|PubMed:8634372}. Note=Associated, through the YDG domain (also called SRA domain), with replicating DNA from early to late S phase, including at replicating pericentric heterochromatin (PubMed:36056023). Also localizes to euchromatic regions. In non-S- phase cells, homogeneously distributed through the nucleus (PubMed:36056023).

Tissue Location

Expressed in thymus, testis, spleen and lung. Within testis, expressed in almost all cells except elongated spermatids.

(Mouse) Uhrf1 Blocking Peptide (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

(Mouse) Uhrf1 Blocking Peptide (C-term) - Images

(Mouse) Uhrf1 Blocking Peptide (C-term) - Background

Multidomain protein that acts as a key epigenetic regulator by bridging DNA methylation and chromatin modification. Specifically recognizes and binds hemimethylated DNA at replication forks via its YDG domain and recruits DNMT1 methyltransferase to ensure faithful propagation of the DNA methylation patterns through DNA replication. In addition to its role in maintenance of DNA methylation, also plays a key role in chromatin modification: through its tudor-like regions and PHD-type zinc fingers, specifically recognizes and binds histone H3 trimethylated at 'Lys-9' (H3K9me3) and unmethylated at 'Arg-2' (H3R2me0), respectively, and recruits chromatin proteins. Enriched in pericentric heterochromatin where it recruits different chromatin modifiers required for this chromatin replication. Also localizes to euchromatic regions where it negatively regulates transcription possibly by impacting DNA methylation and histone modifications. Has E3 ubiquitin-protein ligase activity by mediating the ubiquitination of target proteins such as histone H3 and PML. It is still unclear how E3 ubiquitin-protein ligase activity is related to its role in chromatin in vivo. May be involved in DNA repair.

(Mouse) Uhrf1 Blocking Peptide (C-term) - References

Fujimori A., et al. Mamm. Genome 9:1032-1035(1998).
Davenport J.W., et al. Submitted (JUN-2000) to the EMBL/GenBank/DDBJ databases.
Carninci P., et al. Science 309:1559-1563(2005).
Church D.M., et al. PLoS Biol. 7:E1000112-E1000112(2009).
Muto M., et al. J. Biol. Chem. 277:34549-34555(2002).