

**USP5 Antibody (C-term) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP2134b****Specification**

---

**USP5 Antibody (C-term) Blocking Peptide - Product Information**

Primary Accession [P45974](#)  
Other Accession [UBP5\\_HUMAN](#)

**USP5 Antibody (C-term) Blocking Peptide - Additional Information**

**Gene ID** 8078

**Other Names**

Ubiquitin carboxyl-terminal hydrolase 5, Deubiquitinating enzyme 5, Isopeptidase T, Ubiquitin thioesterase 5, Ubiquitin-specific-processing protease 5, USP5, ISOT

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP2134b](/product/products/AP2134b) was selected from the C-term region of human USP5. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**USP5 Antibody (C-term) Blocking Peptide - Protein Information**

**Name** USP5

**Synonyms** ISOT

**Function**

Deubiquitinating enzyme that participates in a wide range of cellular processes by specifically cleaving isopeptide bonds between ubiquitin and substrate proteins or ubiquitin itself. Affects thereby important cellular signaling pathways such as NF-kappa-B, Wnt/beta-catenin, and cytokine production by regulating ubiquitin-dependent protein degradation. Participates in the activation of the Wnt signaling pathway by promoting FOXM1 deubiquitination and stabilization that induces the recruitment of beta-catenin to Wnt target gene promoter (PubMed: [26912724](http://www.uniprot.org/citations/26912724)). Regulates the assembly and disassembly of heat-induced stress granules by mediating the hydrolysis of

unanchored ubiquitin chains (PubMed:<a href="http://www.uniprot.org/citations/29567855" target="\_blank">29567855</a>). Promotes lipopolysaccharide-induced apoptosis and inflammatory response by stabilizing the TXNIP protein (PubMed:<a href="http://www.uniprot.org/citations/37534934" target="\_blank">37534934</a>). Affects T-cell biology by stabilizing the inhibitory receptor on T-cells PDC1 (PubMed:<a href="http://www.uniprot.org/citations/37208329" target="\_blank">37208329</a>). Acts as a negative regulator of autophagy by regulating ULK1 at both protein and mRNA levels (PubMed:<a href="http://www.uniprot.org/citations/37607937" target="\_blank">37607937</a>). Acts also as a negative regulator of type I interferon production by simultaneously removing both 'Lys-48'-linked unanchored and 'Lys-63'-linked anchored polyubiquitin chains on the transcription factor IRF3 (PubMed:<a href="http://www.uniprot.org/citations/39761299" target="\_blank">39761299</a>). Modulates the stability of DNA mismatch repair protein MLH1 and counteracts the effect of the ubiquitin ligase UBR4 (PubMed:<a href="http://www.uniprot.org/citations/39032648" target="\_blank">39032648</a>). Upon activation by insulin, it gets phosphorylated through mTORC1-mediated phosphorylation to enhance YTHDF1 stability by removing 'Lys-11'-linked polyubiquitination (PubMed:<a href="http://www.uniprot.org/citations/39900921" target="\_blank">39900921</a>). May also deubiquitinate other substrates such as the calcium channel CACNA1H (By similarity).

#### **Cellular Location**

Cytoplasm. Cytoplasm, Stress granule. Nucleus

#### **USP5 Antibody (C-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

#### **USP5 Antibody (C-term) Blocking Peptide - Images**

#### **USP5 Antibody (C-term) Blocking Peptide - Background**

Modification of target proteins by ubiquitin participates in a wide array of biological functions. Proteins destined for degradation or processing via the 26 S proteasome are coupled to multiple copies of ubiquitin. However, attachment of ubiquitin or ubiquitin-related molecules may also result in changes in subcellular distribution or modification of protein activity. An additional level of ubiquitin regulation, deubiquitination, is catalyzed by proteases called deubiquitinating enzymes, which fall into four distinct families. Ubiquitin C-terminal hydrolases, ubiquitin-specific processing proteases (USPs),<sup>1</sup> OTU-domain ubiquitin-aldehyde-binding proteins, and Jab1/Pad1/MPN-domain-containing metallo-enzymes. Among these four families, USPs represent the most widespread and represented deubiquitinating enzymes across evolution. USPs tend to release ubiquitin from a conjugated protein. They display similar catalytic domains containing conserved Cys and His boxes but divergent N-terminal and occasionally C-terminal extensions, which are thought to function in substrate recognition, subcellular localization, and protein-protein interactions.

#### **USP5 Antibody (C-term) Blocking Peptide - References**

Strausberg, R.L., et al., Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903 (2002). Ansari-Lari, M.A., et al., Genome Res. 6(4):314-326 (1996). Falquet, L., et al., FEBS Lett. 376(3):233-237 (1995). Falquet, L., et al., FEBS Lett. 359(1):73-77 (1995).