

USP6 Antibody (N-term) Blocking Peptide
Synthetic peptide
Catalog # BP2135a**Specification**

USP6 Antibody (N-term) Blocking Peptide - Product InformationPrimary Accession
Other Accession[P35125-2](#)
[UBP6_HUMAN](#)**USP6 Antibody (N-term) Blocking Peptide - Additional Information****Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP2135a](/product/products/AP2135a) was selected from the N-term region of human USP6 . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

USP6 Antibody (N-term) Blocking Peptide - Protein Information**USP6 Antibody (N-term) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

USP6 Antibody (N-term) Blocking Peptide - Images**USP6 Antibody (N-term) Blocking Peptide - Background**

Modification of target proteins by ubiquitin participates in a wide array of biological functions. Proteins destined for degradation or processing via the 26 S proteasome are coupled to multiple copies of ubiquitin. However, attachment of ubiquitin or ubiquitin-related molecules may also result in changes in subcellular distribution or modification of protein activity. An additional level of ubiquitin regulation, deubiquitination, is catalyzed by proteases called deubiquitinating enzymes, which fall into four distinct families. Ubiquitin C-terminal hydrolases, ubiquitin-specific processing proteases (USPs),¹ OTU-domain ubiquitin-aldehyde-binding proteins, and Jab1/Pad1/MPN-domain-containing metallo-enzymes. Among these four families, USPs represent the

most widespread and represented deubiquitinating enzymes across evolution. USPs tend to release ubiquitin from a conjugated protein. They display similar catalytic domains containing conserved Cys and His boxes but divergent N-terminal and occasionally C-terminal extensions, which are thought to function in substrate recognition, subcellular localization, and protein-protein interactions.

USP6 Antibody (N-term) Blocking Peptide - References

Paulding, C.A., et al., Proc. Natl. Acad. Sci. U.S.A. 100(5):2507-2511 (2003). Papa, F.R., et al., Nature 366(6453):313-319 (1993). Nakamura, T., et al., Oncogene 7(4):733-741 (1992).