

UNC5B Blocking Peptide (Center)
Synthetic peptide
Catalog # BP21410c**Specification**

UNC5B Blocking Peptide (Center) - Product InformationPrimary Accession [O8IZJ1](#)**UNC5B Blocking Peptide (Center) - Additional Information**

Gene ID 219699

Other Names

Netrin receptor UNC5B, Protein unc-5 homolog 2, Protein unc-5 homolog B, p53-regulated receptor for death and life protein 1, UNC5B, P53RDL1, UNC5H2

Target/Specificity

The synthetic peptide sequence is selected from aa 507-521 of HUMAN UNC5B

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

UNC5B Blocking Peptide (Center) - Protein Information

Name UNC5B

Function

Receptor for netrin required for axon guidance. Mediates axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding. Axon repulsion in growth cones may be caused by its association with DCC that may trigger signaling for repulsion (By similarity). Functions as a netrin receptor that negatively regulates vascular branching during angiogenesis. Mediates retraction of tip cell filopodia on endothelial growth cones in response to netrin (By similarity). It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand (PubMed:12598906). Mediates apoptosis by activating DAPK1. In the absence of NTN1, activates DAPK1 by reducing its autoinhibitory phosphorylation at Ser-308 thereby increasing its catalytic activity (By similarity).

Cellular LocationCell membrane; Single-pass type I membrane protein {ECO:0000250|UniProtKB:O08722}
Membrane raft {ECO:0000250|UniProtKB:O08722}. Note=Associated with lipid rafts.

{ECO:0000250|UniProtKB:O08722}

Tissue Location

Highly expressed in brain. Also expressed at lower level in developing lung, cartilage, kidney and hematopoietic and immune tissues.

UNC5B Blocking Peptide (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

UNC5B Blocking Peptide (Center) - Images**UNC5B Blocking Peptide (Center) - Background**

Receptor for netrin required for axon guidance. Mediates axon repulsion of neuronal growth cones in the developing nervous system upon ligand binding. Axon repulsion in growth cones may be caused by its association with DCC that may trigger signaling for repulsion. It also acts as a dependence receptor required for apoptosis induction when not associated with netrin ligand. Mediates apoptosis by activating DAPK1. In the absence of NTN1, activates DAPK1 by reducing its autoinhibitory phosphorylation at Ser-308 thereby increasing its catalytic activity.

UNC5B Blocking Peptide (Center) - References

Komatsuzaki K., et al. *Biochem. Biophys. Res. Commun.* 297:898-905(2002).
Tanikawa C., et al. *Nat. Cell Biol.* 5:216-223(2003).
Clark H.F., et al. *Genome Res.* 13:2265-2270(2003).
Ota T., et al. *Nat. Genet.* 36:40-45(2004).
Deloukas P., et al. *Nature* 429:375-381(2004).