

**USP13 Antibody (Center) Blocking Peptide**  
**Synthetic peptide**  
**Catalog # BP2141c****Specification**

---

**USP13 Antibody (Center) Blocking Peptide - Product Information**

Primary Accession [O92995](#)  
Other Accession [NP\\_003931](#)

**USP13 Antibody (Center) Blocking Peptide - Additional Information**

**Gene ID** 8975

**Other Names**

Ubiquitin carboxyl-terminal hydrolase 13, Deubiquitinating enzyme 13, Isopeptidase T-3, ISOT-3, Ubiquitin thioesterase 13, Ubiquitin-specific-processing protease 13, USP13, ISOT3

**Target/Specificity**

The synthetic peptide sequence used to generate the antibody [AP2141c](/product/products/AP2141c) was selected from the Center region of human USP13. A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

**Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

**Storage**

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

**Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

**USP13 Antibody (Center) Blocking Peptide - Protein Information**

**Name** USP13

**Synonyms** ISOT3

**Function**

Deubiquitinase that mediates deubiquitination of target proteins such as BECN1, MITF, SKP2 and USP10 and is involved in various processes such as autophagy, endoplasmic reticulum-associated degradation (ERAD), cell cycle progression or DNA damage response (PubMed: [21571647](http://www.uniprot.org/citations/21571647), PubMed: [32772043](http://www.uniprot.org/citations/32772043), PubMed: [33592542](http://www.uniprot.org/citations/33592542)). Component of a regulatory loop that controls autophagy and p53/TP53 levels: mediates deubiquitination of BECN1, a key regulator of autophagy, leading to stabilize the PIK3C3/VPS34-containing complexes.

Alternatively, forms with NEDD4 a deubiquitination complex, which subsequently stabilizes VPS34 to promote autophagy (PubMed:<a href="http://www.uniprot.org/citations/32101753" target="\_blank">32101753</a>). Also deubiquitinates USP10, an essential regulator of p53/TP53 stability. In turn, PIK3C3/VPS34-containing complexes regulate USP13 stability, suggesting the existence of a regulatory system by which PIK3C3/VPS34-containing complexes regulate p53/TP53 protein levels via USP10 and USP13. Recruited by nuclear UFD1 and mediates deubiquitination of SKP2, thereby regulating endoplasmic reticulum-associated degradation (ERAD). Also regulates ERAD through the deubiquitination of UBL4A a component of the BAG6/BAT3 complex. Mediates stabilization of SIAH2 independently of deubiquitinase activity: binds ubiquitinated SIAH2 and acts by impairing SIAH2 autoubiquitination. Regulates the cell cycle progression by stabilizing cell cycle proteins such as SKP2 and AURKB (PubMed:<a href="http://www.uniprot.org/citations/32772043" target="\_blank">32772043</a>). In addition, plays an important role in maintaining genomic stability and in DNA replication checkpoint activation via regulation of RAP80 and TOPBP1 (PubMed:<a href="http://www.uniprot.org/citations/33592542" target="\_blank">33592542</a>). Deubiquitinates the multifunctional protein HMGB1 and subsequently drives its nucleocytoplasmic localization and its secretion (PubMed:<a href="http://www.uniprot.org/citations/36585612" target="\_blank">36585612</a>). Positively regulates type I and type II interferon signalings by deubiquitinating STAT1 but negatively regulates antiviral response by deubiquitinating STING1 (PubMed:<a href="http://www.uniprot.org/citations/23940278" target="\_blank">23940278</a>, PubMed:<a href="http://www.uniprot.org/citations/28534493" target="\_blank">28534493</a>).

#### **Cellular Location**

Cytoplasm.

#### **Tissue Location**

Highly expressed in ovary and testes.

### **USP13 Antibody (Center) Blocking Peptide - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

### **USP13 Antibody (Center) Blocking Peptide - Images**

### **USP13 Antibody (Center) Blocking Peptide - Background**

Modification of target proteins by ubiquitin participates in a wide array of biological functions. Proteins destined for degradation or processing via the 26 S proteasome are coupled to multiple copies of ubiquitin. However, attachment of ubiquitin or ubiquitin-related molecules may also result in changes in subcellular distribution or modification of protein activity. An additional level of ubiquitin regulation, deubiquitination, is catalyzed by proteases called deubiquitinating enzymes, which fall into four distinct families. Ubiquitin C-terminal hydrolases, ubiquitin-specific processing proteases (USPs),<sup>1</sup> OTU-domain ubiquitin-aldehyde-binding proteins, and Jab1/Pad1/MPN-domain-containing metallo-enzymes. Among these four families, USPs represent the most widespread and represented deubiquitinating enzymes across evolution. USPs tend to release ubiquitin from a conjugated protein. They display similar catalytic domains containing conserved Cys and His boxes but divergent N-terminal and occasionally C-terminal extensions, which are thought to function in substrate recognition, subcellular localization, and protein-protein interactions.

### **USP13 Antibody (Center) Blocking Peptide - References**

Puente, X.S., et al., Nat. Rev. Genet. 4(7):544-558 (2003).D'Andrea, A., et al., Crit. Rev. Biochem. Mol. Biol. 33(5):337-352 (1998).Timms, K.M., et al., Gene 217 (1-2), 101-106 (1998).