

AQPEP Blocking Peptide (C-Term)

Synthetic peptide

Catalog # BP21461b

Specification

AQPEP Blocking Peptide (C-Term) - Product Information

Primary Accession

[Q6Q4G3](#)**AQPEP Blocking Peptide (C-Term) - Additional Information**

Gene ID 206338

Other Names

Aminopeptidase Q, AP-Q, 3411-, CHL2 antigen, Laeverin, AQPEP, LVRN

Target/Specificity

The synthetic peptide sequence is selected from aa 844-857 of HUMAN AQPEP

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

AQPEP Blocking Peptide (C-Term) - Protein InformationName LVRN ([HGNC:26904](#))**Function**

Metalloprotease which may be important for placentation by regulating biological activity of key peptides at the embryo-maternal interface. On synthetic substrates it shows a marked preference for Leu-4-methylcoumaryl-7-amide (Leu-MCA) over Met-MCA, Arg-LCA and Lys- LCA. Cleaves the N-terminal amino acid of several peptides such as angiotensin-3, kisspeptin-10 and endokinin C.

Cellular Location

Membrane; Single- pass type II membrane protein

Tissue Location

Specifically expressed in placenta and not in other tissues. Mainly found at the cell surface region of the extravillous trophoblasts. Detected on extravillous trophoblasts in the outer layer of the chorion laeve in the fetal membrane Not detected on either fetal amnionic epithelial cells or maternal decidual cells. Also detected in the migrating extravillous trophoblasts in the maternal decidual tissues (at protein level).

AQPEP Blocking Peptide (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

AQPEP Blocking Peptide (C-Term) - Images

AQPEP Blocking Peptide (C-Term) - Background

Metalloprotease which may be important for placentation by regulating biological activity of key peptides at the embryo- maternal interface. On synthetic substrates it shows a marked preference for Leu-4-methylcoumaryl-7-amide (Leu-MCA) over Met- MCA, Arg-LCA and Lys-LCA. Cleaves the N-terminal amino acid of several peptides such as angiotensin-3, kisspeptin-10 and endokinin C.

AQPEP Blocking Peptide (C-Term) - References

Fujiwara H.,et al.Biochem. Biophys. Res. Commun. 313:962-968(2004).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Schmutz J.,et al.Nature 431:268-274(2004).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Maruyama M.,et al.J. Biol. Chem. 282:20088-20096(2007).