

SLA2 Blocking Peptide (Center)
Synthetic peptide
Catalog # BP21501c**Specification**

SLA2 Blocking Peptide (Center) - Product InformationPrimary Accession [Q9H6Q3](#)**SLA2 Blocking Peptide (Center) - Additional Information****Gene ID** 84174**Other Names**

Src-like-adapter 2, Modulator of antigen receptor signaling, MARS, Src-like adapter protein 2, SLAP-2, SLA2, C20orf156, SLAP2

Target/Specificity

The synthetic peptide sequence is selected from aa 138-149 of HUMAN SLA2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

SLA2 Blocking Peptide (Center) - Protein Information**Name** SLA2**Synonyms** C20orf156, SLAP2**Function**

Adapter protein, which negatively regulates T-cell receptor (TCR) signaling. Inhibits T-cell antigen-receptor induced activation of nuclear factor of activated T-cells. May act by linking signaling proteins such as ZAP70 with CBL, leading to a CBL dependent degradation of signaling proteins.

Cellular Location

Cytoplasm. [Isoform 2]: Cytoplasm. Note=May be cytoplasmic and is not localized to membranes

Tissue Location

Predominantly expressed in immune system, with highest levels in peripheral blood leukocytes. Expressed in spleen, thymus and lymph nodes. Expressed in T-cells as well as in monocytes, and at low level in B-cells. Also detected in placenta, prostate, skin, retina and colon

SLA2 Blocking Peptide (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

SLA2 Blocking Peptide (Center) - Images

SLA2 Blocking Peptide (Center) - Background

Adapter protein, which negatively regulates T-cell receptor (TCR) signaling. Inhibits T-cell antigen-receptor induced activation of nuclear factor of activated T-cells. May act by linking signaling proteins such as ZAP70 with CBL, leading to a CBL dependent degradation of signaling proteins.

SLA2 Blocking Peptide (Center) - References

Holland S.J.,et al.J. Exp. Med. 194:1263-1276(2001).
Loreto M.P.,et al.Oncogene 22:266-273(2003).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Deloukas P.,et al.Nature 414:865-871(2001).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.