

## **UBE2M Antibody (N-term K11) Blocking Peptide**

Synthetic peptide Catalog # BP2169a

## **Specification**

## **UBE2M Antibody (N-term K11) Blocking Peptide - Product Information**

Primary Accession P61081
Other Accession NP 003960

## UBE2M Antibody (N-term K11) Blocking Peptide - Additional Information

**Gene ID 9040** 

#### **Other Names**

NEDD8-conjugating enzyme Ubc12, 632-, NEDD8 carrier protein, NEDD8 protein ligase, Ubiquitin-conjugating enzyme E2 M, UBE2M, UBC12

#### **Target/Specificity**

The synthetic peptide sequence used to generate the antibody <a href=/product/products/AP2169a>AP2169a</a> was selected from the N-term region of human UBE2M . A 10 to 100 fold molar excess to antibody is recommended. Precise conditions should be optimized for a particular assay.

#### **Format**

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

#### Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions**

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## **UBE2M Antibody (N-term K11) Blocking Peptide - Protein Information**

Name UBE2M

Synonyms UBC12

#### **Function**

Accepts the ubiquitin-like protein NEDD8 from the UBA3-NAE1 E1 complex and catalyzes its covalent attachment to other proteins. The specific interaction with the E3 ubiquitin ligase RBX1, but not RBX2, suggests that the RBX1-UBE2M complex neddylates specific target proteins, such as CUL1, CUL2, CUL3 and CUL4. Involved in cell proliferation.

## **UBE2M Antibody (N-term K11) Blocking Peptide - Protocols**



Provided below are standard protocols that you may find useful for product applications.

### • Blocking Peptides

## **UBE2M Antibody (N-term K11) Blocking Peptide - Images**

## **UBE2M Antibody (N-term K11) Blocking Peptide - Background**

The modification of proteins with ubiquitin is an important cellular mechanism for targeting abnormal or short-lived proteins for degradation. Ubiquitination involves at least three classes of enzymes: ubiquitin-activating enzymes, or E1s, ubiquitin-conjugating enzymes, or E2s, and ubiquitin-protein ligases, or E3s. UBE2M is a member of the E2 ubiquitin-conjugating enzyme family. The encoded protein is linked with a ubiquitin-like protein, NEDD8, which can be conjugated to cellular proteins, such as Cdc53/culin.

# **UBE2M Antibody (N-term K11) Blocking Peptide - References**

Gong, L., et al., J. Biol. Chem. 274(17):12036-12042 (1999).Osaka, F., et al., Genes Dev. 12(15):2263-2268 (1998).