

VAC14 Blocking Peptide (N-Term)

Synthetic peptide Catalog # BP21773a

Specification

VAC14 Blocking Peptide (N-Term) - Product Information

Primary Accession

Q08AM6

VAC14 Blocking Peptide (N-Term) - Additional Information

Gene ID 55697

Other Names

Protein VAC14 homolog, Tax1-binding protein 2, VAC14, TAX1BP2, TRX

Target/Specificity

The synthetic peptide sequence is selected from aa 145-159 of HUMAN VAC14

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

VAC14 Blocking Peptide (N-Term) - Protein Information

Name VAC14

Synonyms TAX1BP2, TRX

Function

Scaffold protein component of the PI(3,5)P2 regulatory complex which regulates both the synthesis and turnover of phosphatidylinositol 3,5-bisphosphate (PtdIns(3,5)P2). Pentamerizes into a star-shaped structure and nucleates the assembly of the complex. The pentamer binds a single copy each of PIKFYVE and FIG4 and coordinates both PIKfyve kinase activity and FIG4 phosphatase activity, being required to maintain normal levels of phosphatidylinositol 3- phosphate (PtdIns(3)P) and phosphatidylinositol 5-phosphate (PtdIns(5)P) (PubMed:33098764). Plays a role in the biogenesis of endosome carrier vesicles (ECV) / multivesicular bodies (MVB) transport intermediates from early endosomes.

Cellular Location

Endosome membrane. Microsome membrane {ECO:0000250|UniProtKB:Q80W92}. Note=Mainly associated with membranes of the late endocytic pathway



Tissue LocationUbiquitously expressed.

VAC14 Blocking Peptide (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

• Blocking Peptides

VAC14 Blocking Peptide (N-Term) - Images

VAC14 Blocking Peptide (N-Term) - Background

The PI(3,5)P2 regulatory complex regulates both the synthesis and turnover of phosphatidylinositol 3,5-bisphosphate (PtdIns(3,5)P2). Acts as a positive activator of PIKfyve kinase activity. Also required to maintain normal levels of phosphatidylinositol 3-phosphate (PtdIns(3)P) and phosphatidylinositol 5-phosphate (PtdIns(5)P). Plays a role in the biogenesis of endosome carrier vesicles (ECV) / multivesicular bodies (MVB) transport intermediates from early endosomes.

VAC14 Blocking Peptide (N-Term) - References

Ota T.,et al.Nat. Genet. 36:40-45(2004).

Martin J.,et al.Nature 432:988-994(2004).

Mireskandari A.,et al.Biochim. Biophys. Acta 1306:9-13(1996).

Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

Sbrissa D.,et al.Mol. Cell. Biol. 24:10437-10447(2004).