

CXXC5 Blocking Peptide (N-Term)
Synthetic peptide
Catalog # BP21897a**Specification**

CXXC5 Blocking Peptide (N-Term) - Product Information

Primary Accession [O7LFL8](#)
Other Accession [Q32LB3](#), [Q91WA4](#), [Q5R7N4](#), [Q5XIQ3](#)

CXXC5 Blocking Peptide (N-Term) - Additional Information

Gene ID 51523

Other Names

CXXC-type zinc finger protein 5, CF5, Putative MAPK-activating protein PM08, Putative NF-kappa-B-activating protein 102, Retinoid-inducible nuclear factor, RINF, CXXC5

Target/Specificity

The synthetic peptide sequence is selected from aa 67-80 of HUMAN CXXC5

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

CXXC5 Blocking Peptide (N-Term) - Protein Information

Name CXXC5

Function

May indirectly participate in activation of the NF-kappa-B and MAPK pathways. Acts as a mediator of BMP4-mediated modulation of canonical Wnt signaling activity in neural stem cells (By similarity). Required for DNA damage-induced ATM phosphorylation, p53 activation and cell cycle arrest. Involved in myelopoiesis. Transcription factor. Binds to the oxygen responsive element of COX4I2 and represses its transcription under hypoxia conditions (4% oxygen), as well as normoxia conditions (20% oxygen) (PubMed:23303788). May repress COX4I2 transactivation induced by CHCHD2 and RBPJ (PubMed:23303788). Binds preferentially to DNA containing cytidine-phosphate-guanosine (CpG) dinucleotides over CpH (H=A, T, and C), hemimethylated-CpG and hemimethylated-hydroxymethyl-CpG (PubMed:29276034).

Cellular Location

Nucleus. Cytoplasm {ECO:0000250|UniProtKB:Q5XIQ3} Note=Colocalizes with DVL1 in large bodies localized just outside the nuclear membrane. {ECO:0000250|UniProtKB:Q5XIQ3}

CXXC5 Blocking Peptide (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

CXXC5 Blocking Peptide (N-Term) - Images**CXXC5 Blocking Peptide (N-Term) - Background**

May indirectly participate in activation of the NF- kappa-B and MAPK pathways. Acts as a mediator of BMP4-mediated modulation of canonical Wnt signaling activity in neural stem cells (By similarity). Required for DNA damage-induced ATM phosphorylation, p53 activation and cell cycle arrest. Involved in myelopoiesis.

CXXC5 Blocking Peptide (N-Term) - References

Zhang M.,et al.Sci. China, Ser. C, Life Sci. 52:528-538(2009).
Matsuda A.,et al.Oncogene 22:3307-3318(2003).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Zhou J.,et al.Submitted (JUL-2000) to the EMBL/GenBank/DDBJ databases.
Schmutz J.,et al.Nature 431:268-274(2004).