

LITAF Blocking Peptide (N-Term)
Synthetic peptide
Catalog # BP21944a**Specification****LITAF Blocking Peptide (N-Term) - Product Information**

Primary Accession
Other Accession

[Q99732](#)
[Q9JLJ0](#), [POCOT0](#)

LITAF Blocking Peptide (N-Term) - Additional Information**Gene ID** 9516**Other Names**

Lipopolysaccharide-induced tumor necrosis factor-alpha factor, LPS-induced TNF-alpha factor, Small integral membrane protein of lysosome/late endosome, p53-induced gene 7 protein, LITAF, PIG7, SIMPLE

Target/Specificity

The synthetic peptide sequence is selected from aa 51-60 of HUMAN LITAF

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

LITAF Blocking Peptide (N-Term) - Protein Information**Name** LITAF**Function**

Plays a role in endosomal protein trafficking and in targeting proteins for lysosomal degradation (PubMed:[23166352](http://www.uniprot.org/citations/23166352)). Plays a role in targeting endocytosed EGFR and ERGG3 for lysosomal degradation, and thereby helps down-regulate downstream signaling cascades (PubMed:[23166352](http://www.uniprot.org/citations/23166352)). Helps recruit the ESCRT complex components TSG101, HGS and STAM to cytoplasmic membranes (PubMed:[23166352](http://www.uniprot.org/citations/23166352)). Probably plays a role in regulating protein degradation via its interaction with NEDD4 (PubMed:[15776429](http://www.uniprot.org/citations/15776429)). May also contribute to the regulation of gene expression in the nucleus (PubMed:[10200294](http://www.uniprot.org/citations/10200294), PubMed:[15793005](http://www.uniprot.org/citations/15793005)). Binds DNA

(in vitro) and may play a synergistic role with STAT6 in the nucleus in regulating the expression of various cytokines (PubMed:15793005). May regulate the expression of numerous cytokines, such as TNF, CCL2, CCL5, CXCL1, IL1A and IL10 (PubMed:10200294, PubMed:15793005).

Cellular Location

Cytoplasm. Nucleus. Lysosome membrane; Peripheral membrane protein; Cytoplasmic side. Early endosome membrane. Late endosome membrane. Endosome membrane; Peripheral membrane protein; Cytoplasmic side. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Golgi apparatus membrane. Note=Associated with membranes of lysosomes, early and late endosomes (PubMed:11274176, PubMed:27582497, PubMed:27927196). Can translocate from the cytoplasm into the nucleus (PubMed:15793005). Detected at Schmidt-Lanterman incisures and in nodal regions of myelinating Schwann cells (By similarity) {ECO:0000250|UniProtKB:Q9JLJ0, ECO:0000269|PubMed:11274176, ECO:0000269|PubMed:15793005, ECO:0000269|PubMed:27582497, ECO:0000269|PubMed:27927196}

Tissue Location

Ubiquitously and abundantly expressed. Expressed predominantly in the placenta, peripheral blood leukocytes, lymph nodes and spleen.

LITAF Blocking Peptide (N-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

LITAF Blocking Peptide (N-Term) - Images

LITAF Blocking Peptide (N-Term) - Background

Probable role in regulating transcription of specific genes. May regulate through NFkB1 the expression of the CCL2/MCP-1 chemokine. May play a role in tumor necrosis factor alpha (TNF-alpha) gene expression.

LITAF Blocking Peptide (N-Term) - References

- Polyak K.,et al.Nature 389:300-306(1997).
Myokai F.,et al.Proc. Natl. Acad. Sci. U.S.A. 96:4518-4523(1999).
Moriwaki Y.,et al.J. Biol. Chem. 276:23065-23076(2001).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Bechtel S.,et al.BMC Genomics 8:399-399(2007).