

PDE2A Blocking Peptide (C-Term) Synthetic peptide Catalog # BP22001b

## Specification

# PDE2A Blocking Peptide (C-Term) - Product Information

Primary Accession Other Accession <u>000408</u> <u>P14099, 092254, 001062</u>

## PDE2A Blocking Peptide (C-Term) - Additional Information

Gene ID 5138

**Other Names** cGMP-dependent 3', 5'-cyclic phosphodiesterase, 3.1.4.17, Cyclic GMP-stimulated phosphodiesterase, CGS-PDE, cGSPDE, PDE2A

**Target/Specificity** The synthetic peptide sequence is selected from aa 807-821 of HUMAN PDE2A

#### Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

#### **Precautions** This product is for research use only. Not for use in diagnostic or therapeutic procedures.

## PDE2A Blocking Peptide (C-Term) - Protein Information

Name PDE2A (<u>HGNC:8777</u>)

#### Function

cGMP-activated cyclic nucleotide phosphodiesterase with a dual-specificity for the second messengers cAMP and cGMP, which are key regulators of many important physiological processes (PubMed:<a href="http://www.uniprot.org/citations/15938621" target="\_blank">15938621</a>, PubMed:<a href="http://www.uniprot.org/citations/29392776" target="\_blank">29392776</a>, PubMed:<a href="http://www.uniprot.org/citations/9210593" target="\_blank">9210593</a>). Has a higher efficiency with cGMP compared to cAMP (PubMed:<a

href="http://www.uniprot.org/citations/15938621" target="\_blank">15938621</a>). Plays a role in cell growth and migration (PubMed:<a href="http://www.uniprot.org/citations/24705027" target="\_blank">24705027" target="\_blank">24705027</a>).

Cellular Location [Isoform PDE2A3]: Cell membrane; Lipid- anchor [Isoform PDE2A1]: Cytoplasm



## **Tissue Location**

Widely expressed (PubMed:29392776, PubMed:9210593). Expressed in brain, with high expression observed in the corpus striatum (PubMed:29392776). Also expressed in heart, placenta, lung, skeletal muscle, kidney and pancreas (PubMed:29392776, PubMed:9210593)

# PDE2A Blocking Peptide (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

# Blocking Peptides

PDE2A Blocking Peptide (C-Term) - Images

## PDE2A Blocking Peptide (C-Term) - Background

Cyclic nucleotide phosphodiesterase with a dual- specificity for the second messengers cAMP and cGMP, which are key regulators of many important physiological processes.

## PDE2A Blocking Peptide (C-Term) - References

Rosman G.J., et al.Gene 191:89-95(1997). Bugaj-Gaweda B., et al.Submitted (DEC-2003) to the EMBL/GenBank/DDBJ databases. Ota T., et al.Nat. Genet. 36:40-45(2004). Taylor T.D., et al.Nature 440:497-500(2006). Russwurm C., et al.J. Biol. Chem. 284:25782-25790(2009).