

FLOT2 Blocking Peptide (C-Term)

Synthetic peptide

Catalog # BP22149b

Specification

FLOT2 Blocking Peptide (C-Term) - Product Information

Primary Accession

[Q14254](#)

Other Accession

[A6QLR4](#), [Q60634](#), [Q9Z2S9](#)**FLOT2 Blocking Peptide (C-Term) - Additional Information****Gene ID** 2319**Other Names**

Flotillin-2, Epidermal surface antigen, ESA, Membrane component chromosome 17 surface marker 1, FLOT2, ESA1, M17S1

Target/Specificity

The synthetic peptide sequence is selected from aa 342-356 of HUMAN FLOT2

Format

Peptides are lyophilized in a solid powder format. Peptides can be reconstituted in solution using the appropriate buffer as needed.

Storage

Maintain refrigerated at 2-8°C for up to 6 months. For long term storage store at -20°C.

Precautions

This product is for research use only. Not for use in diagnostic or therapeutic procedures.

FLOT2 Blocking Peptide (C-Term) - Protein Information**Name** FLOT2**Synonyms** ESA1, M17S1**Function**

May act as a scaffolding protein within caveolar membranes, functionally participating in formation of caveolae or caveolae-like vesicles. May be involved in epidermal cell adhesion and epidermal structure and function.

Cellular Location

Cell membrane; Peripheral membrane protein. Membrane, caveola; Peripheral membrane protein. Endosome Membrane; Lipid-anchor. Note=Membrane-associated protein of caveolae

Tissue Location

In skin, expressed in epidermis and epidermal appendages but not in dermis. Expressed in all layers of the epidermis except the basal layer. In hair follicles, expressed in the suprabasal layer

but not the basal layer. Also expressed in melanoma and carcinoma cell lines, fibroblasts and foreskin melanocytes

FLOT2 Blocking Peptide (C-Term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Blocking Peptides](#)

FLOT2 Blocking Peptide (C-Term) - Images

FLOT2 Blocking Peptide (C-Term) - Background

May act as a scaffolding protein within caveolar membranes, functionally participating in formation of caveolae or caveolae-like vesicles. May be involved in epidermal cell adhesion and epidermal structure and function.

FLOT2 Blocking Peptide (C-Term) - References

Zody M.C.,et al.Nature 440:1045-1049(2006).
Schroeder W.T.,et al.J. Biol. Chem. 269:19983-19991(1994).
Kalnine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.
Gorbea C.,et al.J. Biol. Chem. 285:31616-31633(2010).
Burkard T.R.,et al.BMC Syst. Biol. 5:17-17(2011).